

THREE-DIMENSIONAL ANALYSIS OF PALATE ROUGHNESS AS A BIOMETRIC IDENTIFICATION TOOL

Monique Miranda Araújo¹, Xaiane de Oliveira², Ricardo Almeida Caribé de Freitas³
Jeidson Antônio Morais Marques⁴, Jamilly de Oliveira Musse⁵

Highlights:

(1) The three-dimensional analysis of palatal roughness is a methodological proposal applicable to human identification processes.

(2) The three techniques used in the analysis of palatal rugae (Silva - 1938), Carrea - 1937 and Bassauri - 1961) were not sufficient for human identification when used in isolation.

(3) The individual code generated by the association of the three methods proved to be unique for each model used, proving the applicability of the three-dimensional analysis of palatal wrinkles for human identification.

PRE-PROOF

(as accepted)

This is a preliminary and unedited version of a manuscript that has been accepted for publication in Context and Health Journal. As a service to our readers, we are providing this initial version of the manuscript as accepted.

The manuscript will still undergo revision, formatting, and approval by the authors
before being published in its final form

<http://dx.doi.org/10.21527/2176-7114.2024.48.13525>

How to cite:

Araújo MM, de Oliveira X, de Freitas RAC, Marques JAM, Musse J de O. Three-dimensional analysis of palate roughness as a biometric identification tool . Rev. Contexto & Saúde, 2024;24(48): e13525

1 Unidade de Ensino Superior de Feira de Santana – UNEF. Feira de Santana/BA, Brasil.

<https://orcid.org/0009-0002-0229-5829>

2 Unidade de Ensino Superior de Feira de Santana – UNEF. Feira de Santana/BA, Brasil.

<https://orcid.org/0009-0000-8063-8992>

3 Unidade de Ensino Superior de Feira de Santana – UNEF. Feira de Santana/BA, Brasil.

<https://orcid.org/0009-0005-3042-6402>

4 Unidade de Ensino Superior de Feira de Santana – UNEF. Feira de Santana/BA, Brasil.

<https://orcid.org/0000-0003-3070-7077>

5 Unidade de Ensino Superior de Feira de Santana – UNEF. Feira de Santana/BA, Brasil.

<https://orcid.org/0000-0001-5769-9228>

THREE-DIMENSIONAL ANALYSIS OF PALATE ROUGHNESS
AS A BIOMETRIC IDENTIFICATION TOOL

ABSTRACT

Palatal wrinkles are elevations of the oral mucosa that cover the palatine process and are located on the maxillary bone. Because its anatomy is immutable and unique to each individual, the method of human identification using palatal wrinkles has a lot of potential to become indisputable. Currently, with the incorporation of new technologies in dentistry, three-dimensional analysis of palatal wrinkles using scans has become a reality. The aim of this study was to analyze the applicability of three-dimensional analysis of palatal wrinkles for human identification, through the observation of 3D images resulting from digital scanning. This is an experimental study with a qualitative-quantitative approach, whose sample consisted of the scanning of 50 plaster models, where three-dimensional images were generated, the classifications used were with the techniques of Silva (1938), Carrea, (1937) and Bassauri (1961). The 3D scanning was done by a software system from the Zirkonzahn company, using the s600 arti scanner, and the wrinkle analysis was carried out by the Meshmixer program. When used separately, the techniques were not sufficient for individual human identification. However, the individual code generated by combining the three methods proved to be unique for each model used. Thus, the results obtained proved the applicability of three-dimensional analysis of palatal wrinkles for human identification.

KEYWORDS: Human Identification, Hard Palate, Forensic Odontology

1. INTRODUCTION

The purpose of human identification is to reveal the identity of a body. It can be carried out on both living and post-mortem individuals, in a wide variety of conditions (preserved, decomposing, charred, dismembered, mutilated and fragmented). It is carried out by trained technicians or professionals with differentiated and specific knowledge, and consists of comparing characters, looking for coincidences between previously recorded data and that obtained in the present¹.

In forensic anthropology, one of the most difficult tasks is human identification, which can be carried out using fingerprints, DNA comparisons and dental information².

The materials currently used to apply the dental arch identification technique are the documents produced during dental care, which generally consist of radiographs, dental records, plaster models and photographs³, and the examination is carried out in three stages: examination of the cadaver's dental arches, examination of the suspect's dental documents and comparison, with the dental expert reaching a conclusion about the individual's identity. If plaster models or digitized images of palatal rugoscopy are available, their detailed analysis can also be used as a parameter for comparison, since palatal wrinkles are unique and do not change even after morphological alterations, orthodontic and surgical treatments in the palate region, and can resist decomposition after death for up to seven days¹.

Due to the immutable and unique anatomy of each individual, the method of human identification using palatal wrinkles has the potential to become indisputable. Its study as a method of individual identification was first mentioned by Lopes de Léon in 1924, inspired by Vucetich. From then on, research began into palatal wrinkles and the quickest and safest ways of identifying people using them, with various classification systems being proposed, such as Carrea (1937), Silva (1938), Martins dos Santos (1952), Bassauri (1961), Thomas and Kotze (1983), among others⁴⁻⁵.

Currently, with the incorporation of new technologies in dentistry, the imaging of dental arches from intraoral scanning has become a reality⁶.

According to Souza (2020)⁷, With the use of scanning, it is possible to reproduce the dental arch and analyze it in an average of 10 minutes. It is a fast and reliable method, which normally does not require repetition, providing a safe method of comparing the arches to be analyzed, since it does not alter the shape like traditional molding.

With the increasing use of technology, digital dentistry is a reality and is being used more and more⁵. Considering its possible applicability for human identification through palatal wrinkles, this study aimed to analyze the applicability of three-dimensional analysis of palatal wrinkles for human identification, based on images resulting from 3D scanning.

3. METHODOLOGY

This is an experimental study with a qualitative and quantitative approach, which analyzed the applicability of palatal wrinkles as a method of human identification through three-dimensional images. The methods for classifying palatal wrinkles defined by Carrea (1937)⁸ were adopted as categorical variables, Silva (1938)⁹ and Bassauri (1961)¹⁰.

A pilot study was carried out beforehand in order to calibrate the researchers, test the data collection methods and the program, and this stage was analyzed by calculating the intraclass correlation coefficient.

Data was collected by undergraduate dentistry students from the Feira de Santana Higher Education Unit (UNEF). The data obtained was analyzed by a pair of observer-researchers, who classified the palatal roughness using three-dimensional images.

Fifty plaster models, 25 female and 25 male, were scanned using a software system from the Zirkonzahn company, using the s600 arti scanner, thus generating files in STL format. Data analysis was carried out using Meshmixer, a free software program used for editing 3D meshes (Figure 1).




Figure 1 - Three-dimensional image of palatal wrinkles.

The images of the upper models were made available on a pen drive and analyzed according to the classifications of Carrea (1937)⁸, Silva (1938)⁹ and Bassauri (1961)¹⁰. The data was recorded on a specific form, already validated in a previous study by the researcher responsible for this project, using the plaster model and not the 3D image.

The Carrea Method (1937)⁸, was used to delineate palatal roughness by dividing the wrinkles into right and left, separating them by a vertical line up to the incisive papilla. Based on the bilateral direction, the wrinkles were analyzed under four circumstances: wrinkles that converge on the palatine raphe in the posteroanterior direction were type I, those located horizontally and perpendicular to the palatine raphe were type II and wrinkles that are directed anteroposteriorly and converge on the palatine raphe were type III. Finally, those in different directions were classified as type IV (Figure 1).




Figure 1. Classification proposed by Carrea (1937) Source: FONSECA, FLÓREZ (2009)¹¹.

In Silva's (1938)⁹, method, wrinkles were analyzed individually, adopting the following conformation: type 1: straight; type 2: curved; type 3: angled; type 4: circular; type 5: sinuous, type 6: dotted and compound "Y", cup, racket or branched wrinkles. Wrinkles smaller than two millimeters were disregarded. The method ends with the sum of the wrinkles, according to the quantity of each conformation (Figure 2).




Figure 2. Silva's classification (1938).

Source: CASTRO-SILVA, SILVA E VEIGA (2014)¹².

The Bassauri Method (1961) analyzes the shape of the palatal wrinkles. The main wrinkle, located more anteriorly, was classified with capital letters (A, B, C, D, E, F, X) and the accessory wrinkles with numbers (1, 2, 3, 4, 5, 6, 7), thus allowing classification into more than one type and on both sides of each impression (Chart 2).

Table 2. Classification system proposed by Bassauri.

Main wrinkle	Accessory wrinkle	Anatomy
A	1	Point
B	2	Line
C	3	Angle
D	4	Sinuuous
E	5	Curve
F	6	Circle
X	7	Polymorphic

Source: TORNOVOI; SILVA (2010)¹³.

At the end of the analysis and classification of all the 3D images, an electronic table was drawn up with the information on palatal roughness in the Excel program (Office 2010, Microsoft Corporation, Redmond, Washington, USA), showing the quantitative values of the palatal wrinkles. Three methods were applied with individualized codes in order to prove the uniqueness of the palatal wrinkles.

The study complies with National Health Council Resolution No. 466/12, which regulates ethics in research involving human beings, and was approved by the research ethics committee of the State University of Feira de Santana, under CAAE 37278920.2.0000.0053.

4. RESULTS

Carrea's (1937) method found a predominance of type IV wrinkles (36%), followed by type II (34%) and type III (28%). Type I (2%) was the least common type of wrinkle found in this study's total group.

Table 3. Evaluation of palatal roughness according to Carrea's method (1937), Feira de Santana, Bahia, Brazil, 2021.

Classification Frequency Percentage	Frequency	Percentage
I	1	2,0
II	17	34,0
III	14	28,0
IV	18	36,0
Total	50	100,0

Morfotipos Silva's (1938) rugoscopic classification is based on numerical morphotypes that define palatal roughness (type 1: straight; type 2: curved; type 3: angular; type 4: circular; type 5: sinuous; type 6: point and type 7: compound), with the sum of wrinkles ranging from simple to compound. This type of analysis found values ranging from 16 to 57, with an average of 32.4 (Chart 4).

Table 4. Evaluation of palatal roughness according to Silva's method (1938). Feira de Santana, Bahia, Brazil, 2021.

Measurements	Values
Average	32,4
Minimumm	16
Maximum	57

According to the analysis of palatal wrinkles using the Bassauri (1961) method, shown in Table 5, the wrinkles were classified according to their morphotypes. In this method, two models were classified as LDE44424LEF524522 and two others as LDX2421LED245.

Table 5. Evaluation of palatal roughness according to the Bassauri method (1963), Feira de Santana, Bahia, Brazil, 2021.

Classification	Frequency	Percentage
LDB15554LEE24552	1	2,0
LDB242LEB224	1	2,0
LDB252LEB4455	1	2,0
LDB4445LEX442	1	2,0
LDB44554LEX7432	1	2,0
LDB4544LEB425244	1	2,0
LDB7525LEE422	1	2,0
LDC444LEC44222	1	2,0
LDD224LED442	1	2,0
LDD452LEF74	1	2,0
LDD5445LEE7432	1	2,0
LDE14522LEE7544	1	2,0
LDE2252555LEE444	1	2,0
LDE2414LEE4445	1	2,0
LDE244LED245	1	2,0
LDE24532LEE24423	1	2,0
LDE24544LEF24	1	2,0
LDE25424LEE222	1	2,0
LDE25455LEX145	1	2,0
LDE4144LEX4117	1	2,0
LDE436LEB45152	1	2,0
LDE44424LEF524522	2	4,0
LDE44444LEE4445	1	2,0
LDE45252LEE2545	1	2,0

LDE452LEE5445	1	2,0
LDE542LEE24	1	2,0
LDE5445LEE2534	1	2,0
LDE732LEX44	1	2,0
LDE741LEF444	1	2,0
LDE74LEX4	1	2,0
LDEX754LEX2427	1	2,0
LDF532LEX143	1	2,0
LDX21444ÇLEX24444	1	2,0
LDX22LEF27	1	2,0
LDX2421LED245	2	4,0
LDX2445LEX42122	1	2,0
LDX254LEB545	1	2,0
LDX25LEF545	1	2,0
LDX314LED442	1	2,0
LDX4142LEX457	1	2,0
LDX414LEX5445	1	2,0
LDX434LED7445	1	2,0
LDX44LEX44	1	2,0
LDX454LEX424	1	2,0
LDX4LEE242	1	2,0
LDX52LEF5	1	2,0
LDX52LEX54	1	2,0
LDX5434LEX474	1	2,0
Total	50	100,0

The combination of the three methods generated an individual code (Chart 6) for each model used, proving the uniqueness of the palatal wrinkles.

Table 6. Evaluation of individual code units, Feira de Santana, Bahia, Brazil, 2021.

Code	Frequency	Percentage
I23LDE45252LEE2545	1	2,0
II16LDB242LEB224	1	2,0
II20LDB252LEB4455	1	2,0
II21LDE25424LEE222	1	2,0
II22LDB7525LEE422	1	2,0
II23LDX25LEF545	1	2,0
II25LDX254LEB545	1	2,0
II28LDD224LED442	1	2,0
II30LDE2252555LEE444	1	2,0
II30LDE24544LEF24	1	2,0
II31LDE436LEB45152	1	2,0
II32LDX2421LED245	1	2,0
II33LDB44554LEX7432	1	2,0
II34LDC444LEC44222	1	2,0
II36LDB4445LEX442	1	2,0
II37LDX314LED442	1	2,0
II37LDX454LEX424	1	2,0
II39LDE25455LEX145	1	2,0
II54LDE4144LEX4117	1	2,0
III18LDE542LEE24	1	2,0
III21LDX22LEF27	1	2,0
III21LDX52LEF5	1	2,0
III26LDE452LEE5445	1	2,0
III26LDE74LEX4	1	2,0
III27LDE732LEX44	1	2,0
III29LDD452LEF74	1	2,0
III31LDB15554LEE24552	1	2,0
III35LDD5445LEE7432	1	2,0
III44LDX434LED7445	1	2,0

III45LDX4142LEX457	1	2,0
III46LDX5434LEX474	1	2,0
III47LDX414LEX5445	1	2,0
IV21LDX4LEE242	1	2,0
IV24LDX52LEX54	1	2,0
IV26LDE244LED245	1	2,0
IV29LDE5445LEE2534	1	2,0
IV31LDF532LEX143	1	2,0
IV32LDE741LEF444	1	2,0
IV33LDX2421LED245	1	2,0
IV34LDE24532LEE24423	1	2,0
IV34LDE44424LEF524522	1	2,0
IV34LDX44LEX44	1	2,0
IV35LDE44424LEF524522	1	2,0
IV38LDB4544LEB425244	1	2,0
IV38LDE14522LEE7544	1	2,0
IV38LDE2414LEE4445	1	2,0
IV41LDX2445LEX42122	1	2,0
IV42LDEX754LEX2427	1	2,0
IV46LDE44444LEE4445	1	2,0
IV57LDX21444ÇLEX2444 4	1	2,0
Total	50	100,0

5. DISCUSSION

This study used the method of human identification by palatal wrinkles, respecting their biological and anatomical aspects for an accurate and coherent analysis. According to Brígido (2018)², wrinkles remain stable throughout life, undergoing only a few changes in position and length, triggered by the natural development of the individual.

In view of the diversity of methods for analyzing palatal wrinkles, it is necessary to validate and standardize their use. Thus, for surveys to be reliable, it is essential to have methodological rigor so that the data can be reproduced legitimately and reliably. In this

sense, Andrade (2004)¹⁴ affirms that intra-examiner and extra-examiner agreement is an important process for reducing errors in surveys. Influencing the accuracy of data collection. For this reason, intra-examiner calibration was carried out in this study, obtaining an intraclass correlation coefficient considered by McBride (2005)¹⁵ to be almost perfect, with a value of 0.9985.

A study carried out by Veloso (2017)⁴, using Carrea's method (1937) on palatal wrinkles in monozygotic twins in the city of João Pessoa/PB, found that the twins had between 8 and 14 wrinkles, with type IV wrinkles prevailing. The most common form of wrinkle was the sinuous type.

Castro-Silva, Silva and Veiga (2014)¹² carried out a study in Niterói-RJ, in 2013, with the aim of assessing the effectiveness of palatal rugoscopy as a human identification technique, using 184 plaster models and applying Carrea's (1937) method. It was observed that the individuals had a higher prevalence of type IV. These data were similar to the present study, where there was a higher prevalence of type IV wrinkles, which were found in 36% of the models analyzed, followed by type II (34%). Type I was the least common (2%).

España et al. (2010)¹⁶ carried out a data storage study using the method proposed by Silva (1934) on a sample of 20 firefighters from the University of Los Andes Mérida in Venezuela. The study models were analyzed to create a database with individualized palatal wrinkle characteristics for each of the firefighters analyzed. The study showed an average of 13 wrinkles for each individual. This fact, however, is different to that found in this study, with an average of 32.4 wrinkles per individual, a higher average than the result presented by España.

Arone et al. (2018)¹⁷ analyzed the characteristics of palatal wrinkles at the UNSAAC dental clinic in Cusco, Peru, using the Bassauri (1963) identification technique. An analysis was carried out on 62 study models and intraoral photographs obtained from 28 men and 34 women. The aim of the study was to identify the individuals using Bassauri's classification. Each individual generated a unique identification code, confirming the individuality of the wrinkles. This study, however, showed a different result. In the Bassauri (1963) technique, there were two situations in which the classifications were repeated with the code LDE44424LEF524522 and LDX2421LED245, contradicting the result found by Arone et al. (2018), confirming that the Bassauri technique alone is not sufficient for identification. The combination of the techniques proposed by Carrea (1937), Silva (1938) and Bassauri (1961) generated a unique code for each individual, proving uniqueness.

According to Oliveira (2015)¹⁸ with the current use of digital technology, it is possible to analyze palatal wrinkles through 3D scanning, which gives us precise quality in three-dimensional images. Scanning is a non-invasive, fast, reliable and high-resolution method that makes it possible to obtain a copy of models and the human dental arch in any state. Cardoso et al. (2018)¹⁹ state that, although traditional molding is more commonly used in dental offices, it is known that the plaster model can cause alterations if it is not stored correctly, making the impressions obtained inadequate for a possible comparison of images at the time of human identification.

For this reason, this research opted for 3D scanning, because according to Pagano et al. (2019)²⁰ they state that digital information has brought us a great advantage, making the work faster, reducing the chance of error and generating quality images. In addition, they allow us to easily change the file as many times as necessary to simulate different angles and situations.

Studies are currently being carried out on the use of 3D scanners to assess palatal roughness. According to Oliveira et al. (2015)¹⁸, CAD/CAM has become increasingly used as a tool in dentistry, being used to print these models and has been used for research due to its ease of handling and the quality of the three-dimensional images.

According to Cechelero (2021)²¹ the material used for molding considered to be the gold standard is elastomer and even then it has disadvantages when compared to scanning, such as: distortions related to molding, execution time, material costs, patient discomfort, the need to disinfect the trays and the operator-dependent technique. Digital scanning is a technique that has been reported to be faster and more convenient for both the patient and the dentist, with digital impressions being more accurate and superior to traditional impressions, according to the literature.

Therefore, according to Souza (2020)⁷, three-dimensional images are, in fact, a reality in dentistry and have been used in several areas, including prosthodontics, implantology, dentistry, orthodontics, endodontics, orthognathic surgery, and now in forensic dentistry, helping with human identification through palatal wrinkles.

It's worth noting that although these anatomical structures meet the biological and technical requirements that make them important in forensic investigations, such as stability, durability, uniqueness, practicality and classifiability, the diversity of classification alternatives makes them controversial in terms of their reliability. For this reason, it is suggested that different forms of classification be combined, as was done in this study.

6. CONCLUSION

The three-dimensional analysis of palatine wrinkles is a useful method that can be applied in Official Forensic Institutes. The combination of the three classification methods used in this study proved the uniqueness of palatal wrinkles.

The scanner proved effective in classifying wrinkles, simplifying and facilitating the process. It was possible to move the plaster models 360°, zoom in and zoom out without distortions and alterations, allowing the details of each wrinkle to be visualized and classified according to the proposed methods.

In this way, the technique of human identification through the analysis of palatal wrinkles using 3D scanning becomes an effective method with a lot of potential to be introduced into the routine of forensic odontology

REFERENCES

1. Neves ISR, Santiago APACS, Silva MIT, Oliveira ERM. Rugoscopia palatina e seus desafios na identificação humana: uma revisão de literatura. *Research, Society and development* 2021, 10(5):1-11.
2. Brigídio JA. Rugoscopia Palatina na Odontologia Forense. *Revista Diálogos Acadêmicos* 2018, 7(1):15-19.
3. Silva D, Mercedes L, Andrade E, Castro M. A Importância dos Arcos Dentários e da Rugoscopia Palatina na Identificação Humana. *J Business Techn* 2020, 5(1): 41-52.
4. Veloso CVL. Estudo rugoscópico em gêmeos monozigóticos na cidade de João Pessoa/PB. João Pessoa: Universidade Federal da Paraíba, 2017.
5. Barrera AS, Pacheco RAE, Quispe LRJ. Relevância de las rugas palatinas como método de identificación forense. *Revista Científica Odontol* 2020, 2(8): 89-95.
6. Bósio JÁ, Del Santo M, Jacob HB. Odontologia Digital Contemporânea– Scanners Intraorais Digitais. *Orthod. Sci. Pract* 2017, 10 (39): 355-362.
7. Souza CM. FLUXO DIGITAL: Uma Realidade na Odontologia Reabilitadora. Salvador: Escola de Medicina e Saúde Pública, 2020.
8. Carrea JU. La Identificación Humana Por Las Rugosidades Palatinas. *RevOrthodont* 1937, 1(1): 3-23.
9. Silva L. Ficha rugoscópica palatina. *Brasil Odonto* 1938, 14(1): 307-16.
10. Basauri C. Forensic Odontology and Identification. *Int Crim Police Rev* 1961, 16(1):

45-51.

11. Fonseca GM, Flórez CDR. Propuesta De Codificación Y Análisis De Rugosidades Palatinas Para Su Aplicación Em Odontología Antropológica Y Forense. *Revista Oral* 2009, 10(31): 518-523.
12. Castro-Silva, Silva OML, Veiga BMC. Use Of Palatine Rugoscopy As Biometric Tool: A Population Study In Niteroi, RJ, Brazil. *Rev. Odontol. UNESP* 2014, 3(43):203-208.
13. Tornavoi DC, Silva RHA. Rugoscopia Palatina e a Aplicabilidade na Identificação Humana em Odontologia Legal: Revisão de Literatura. *Ética&Justiça* 2010, 15(1): 28-34, 2010.
14. Andrade FB, Mathias R, Miotto MHMB, Coutinho JS. Importância da calibração de examinadores para estudos epidemiológicos em odontologia. *Vitória* 2004, 6(1): 16-23.
15. McBride GB. A proposal for strength-of-agreement criteria for Lin's Concordance Correlation Coefficient. *NIWA Client Report HAM* 2005, 1(1): 62.
16. España L, Paris A, Florido R, Arteaga F, Solórzano E. Estudio De Las Características Individualizantes De Las Rugas Palatinas. Caso: Bomberos De La Universidad De Los Andes Mérida. Venezuela. *Cuad Med Forense* 2010, 16(4): 199-204.
17. Arone F, Eliceni S. Características Individuales De Las Rugas Palatinas En Estudiantes De La Clínica Odontologica De La Unsaac, Mediante La Clasificación De Basauri, Cusco. Tesis para optar al título profesional de cirujano dentista. Cusco. Universidade Nacional de San Antonio Abad del, 2018.
18. Oliveira GS, MARQUES JAM. Uso da Tecnologia de Impressão Tridimensional na Rugoscopia Palatina. *RBOL* 2015, 2(2):21-34.
19. Cardoso FL, Alberfaro KPA, Ribeiro S, Assis VKS, Reis LO. Moldagem Digital em Odontologia: Perspectivas Frente à Convencional- Uma Revisão de Literatura. *II Jornada de Iniciação Científica da FACIG* 2018, 3(4): 1-6.
20. Pagano S. Evaluation of the Accuracy of Four Digital Methods by linear and Volumetric Analysis of Dental Impressions. *MDPI Journal* 2019; 12(1958): 1- 20.
21. Cechelero EB, Bellan MC, BISI MA. Análise Comparativa de Técnicas de Escaneamento Digital: Estudo In Vitro. *Arch Health Invest* 2021, 10(2): 248-254.

Submitted: 30/7/2022

Accepted: 24/8/2023

Published: 23/2/2024

Authors' Contributions:

Monique Miranda Araújo: Conceptualization; Methodology; Software; Validation; Visualization; Writing – original draft.

Xaiane de Oliveira: Conceptualization; Methodology; Resources.

Ricardo Almeida Caribé de Freitas: Methodology; Software; Validation; Visualization; Writing – original draft.

Jeidson Antônio Moraes Marques: Conceptualization; Data curation; Formal analysis; Supervision; Writing – original draft; Writing – review & editing.

Jamilly de Oliveira Musse: Conceptualization; Data curation; Formal analysis; Project administration; Supervision; Writing – original draft; Writing – review & editing.

All authors have approved the final version of the text.

Conflict of interest: There are no conflicts of interest.

Financing: Does not have financing.

Corresponding Author:

Jamilly de Oliveira Musse

Unidade de Ensino Superior de Feira de Santana – UNEF

Av. Luís, Av. Eduardo Magalhães Subaé - Aviário, Feira de Santana - BA, Brasil. CEP 44079-002

E-mail: jamillymusse@gmail.com

Editor: Dr. Matias Nunes Frizzo

Editor-in-chief: Dra. Adriane Cristina Bernat Kolankiewicz

This is an open access article distributed under the terms of the Creative Commons license.

