

ORIGINAL ARTICLE

Occurrence and Characteristics of Elderly with Covid-19 in Southern Brazil

Luana Schayene Wagner¹; Gabrieli de Fátima Prudêncio dos Santos²
Juliana Lemes dos Santos³; Tatiana Mugnol⁴; Diego Pascoal Golle⁵
Josiane Woutheres Bortolotto⁶; Mariana Migliorini Parisi⁷, Janaina Coser⁸

Highlights:

1. 24.2% of elderly people a positive result in the RT-qPCR test for SARS-CoV-2.
2. Females had a higher proportion of positive cases compared to males.
3. A higher proportion of infected with pre-existing cardiovascular diseases.

ABSTRACT

This study aimed to describe the occurrence of Covid-19 in elderly patients treated at public health services in southern Brazil cities and their clinical and sociodemographic characteristics. This is a cross-sectional, observational, and retrospective study. From July 2020 to May 2021, medical records of elderly patients who underwent a test for Covid-19 at the molecular diagnostics laboratory of the University of Cruz Alta were selected. 310 elderly patients, with an average age of 60 to 99 years, were included in the study. Information was collected regarding the result of the RT-qPCR test, gender, symptoms, and comorbidities. We identified that 24.2% of the elderly evaluated in the study had a positive molecular diagnosis for Covid-19. The average age of the sample studied was 69 years (± 8.89). Females had a higher proportion of positive cases (69.3%) compared to males (30.7%) ($p=0.017$). A higher proportion of individuals infected with Sars-Cov2 among those who reported cough, fever $>37^{\circ}\text{C}$ and pre-existing cardiovascular diseases ($p = 0.028$; 0.015 ; 0.017 respectively) was also identified. Learning the clinical and sociodemographic profile of the elderly with Covid-19 can contribute to strategies for care and prevention of the disease in this group. The study findings indicate that greater attention should be given to the elderly with heart diseases and those who have fevers and coughs.

Keywords: Test for Covid-19; Elderly Health; Comprehensive health care.

OCORRÊNCIA E CARACTERÍSTICAS DE IDOSOS COM COVID-19 NO SUL DO BRASIL

RESUMO

Este estudo teve como objetivo descrever a ocorrência de Covid-19 em pacientes idosos atendidos em serviços públicos de saúde de cidades do Sul do Brasil e as suas características clínicas e sociodemográficas. Trata-se de um estudo transversal, observacional e retrospectivo. Foram selecionados prontuários de pacientes idosos que realizaram teste para Covid-19 no laboratório de diagnóstico molecular da Universidade de Cruz Alta, no período de julho de 2020 a maio de 2021. Foram incluídos no estudo 310 pacientes idosos com idades entre 60 e 99 anos. Foram coletadas informações quanto ao resultado do teste RT-qPCR, sexo, sintomas e comorbidades. Identificamos que 24,2% dos idosos avaliados no estudo tiveram diagnóstico molecular positivo para Covid-19. A média de idade da amostra estudada foi de 69 anos ($\pm 8,89$). O sexo feminino apresentou maior proporção de casos positivos (69,3%) em comparação ao sexo masculino (30,7%) ($p=0,017$). Identificamos uma maior proporção de indivíduos infectados com Sars-Cov2 entre aqueles que relataram tosse, febre $>37^{\circ}\text{C}$ e doenças cardiovasculares preexistentes ($p = 0,028$; $0,015$; $0,017$ respectivamente). Conhecer o perfil clínico e sociodemográfico dos idosos com Covid-19 pode contribuir com estratégias de cuidado e prevenção da doença neste grupo. Os achados do estudo mostram que maior atenção deve ser dada aos idosos com cardiopatia e que apresentam febre e tosse.

Palavras-chave: teste para Covid-19; saúde do idoso; assistência integral à saúde.

¹ Universidade de Cruz Alta – Unicruz. Cruz Alta/RS, Brasil. <https://orcid.org/0000-0002-9763-6296>

² Universidade de Cruz Alta – Unicruz. Cruz Alta/RS, Brasil. <https://orcid.org/0000-0002-9765-6836>

³ Universidade de Cruz Alta – Unicruz. Cruz Alta/RS, Brasil. <https://orcid.org/0000-0003-1724-4477>

⁴ Universidade de Cruz Alta – Unicruz. Cruz Alta/RS, Brasil. <https://orcid.org/0000-0002-1003-5800>

⁵ Faculdade Santo Ângelo (FASA). Santo Ângelo/RS, Brasil. <https://orcid.org/0000-0002-5264-8007>

⁶ Universidade de Cruz Alta – Unicruz. Cruz Alta/RS, Brasil. <https://orcid.org/0000-0003-0271-1685>

⁷ Universidade de Cruz Alta – Unicruz. Cruz Alta/RS, Brasil. <https://orcid.org/0000-0002-2298-7809>

⁸ Universidade de Cruz Alta – Unicruz. Cruz Alta/RS, Brasil. <https://orcid.org/0000-0003-3631-0847>

INTRODUCTION

At the end of 2019, the World Health Organization (WHO) was alerted to the occurrence of a series of severe pneumonia cases, similar to viral pneumonia, which began in the city of Wuhan, China (Huang 2020)¹. Through sequencing analysis of samples from the lower respiratory tract, it was discovered that it was a new, highly pathogenic and transmissible virus, which was identified as Severe Acute Respiratory Syndrome Coronavirus Virus 2 (Sars-CoV-2), responsible for the Coronavirus Disease 2019 (Covid-19)¹⁻².

Due to its high transmissibility, the virus spread quickly and on January 30th, 2020 the WHO declared a public health emergency of international concern and, on March 11th, the Covid-19 pandemic was officially declared²⁻³. Since then, the numbers of cases and deaths related to the disease have increased significantly, showing the high incidence worldwide, with Brazil being one of the most affected countries, reaching the 3rd place in the world ranking⁴.

The transmission of the virus occurs by direct person-to-person contact or by indirect contact through contaminated objects or surfaces^{3,5}. Among the commonly reported symptoms are fever, cough, shortness of breath, anosmia and ageusia. However, some individuals, even those infected, may not have symptoms and still transmit the virus, which makes it difficult to contain the pandemic^{3,6}.

Through studies related to the pathogenesis of this disease, some risk factors associated with higher fatality rates from Covid-19 were considered, such as advanced age, the male gender, obesity, smoking, and presence of comorbidities⁶⁻⁷.

WHO data and the literature indicate that the largest number of confirmed cases in the world is concentrated in the age range between 25 and 64 years, with an average age of 49 years. However, people aged 65 years and over, and/or with pre-existing diseases, had the highest death rates, with about 73% of the total deaths recorded, a situation that places them as one of the main risk groups for Covid-19⁸⁻⁹. The literature justifies age as a risk factor for the disease due to the lower responsiveness of the immune system and the senescence of the respiratory system with advancing age. Thus, considering the tropism of Covid-19 through this system, the elderly become more vulnerable to the respiratory complications of the disease and have greater difficulty in fighting the virus^{6,8}.

In fact, since the beginning of the pandemic, the literature has reported that Sars-CoV-2 infection affects all age groups¹⁰, however, it has also been widely reported that the elderly is more susceptible to Covid-19, and those with underlying diseases, are considered a high-risk group¹¹. Therefore, early diagnosis and individualized therapeutic management must be developed for elderly people, based on their clinical history and presence of comorbidities¹². However, in Brazil, specific studies that investigated the clinical spectrum of Covid 19 in the elderly community population¹³ are rare, covering other research focuses¹⁴ and/or concentrating on in institutionalized elderly¹⁵⁻¹⁹, a context similar to studies developed in the southern region of Brazil²⁰⁻²².

In view of the above, the present study sought to analyze the occurrence and characteristics of Covid-19 in an elderly population (elderly peoples aged 60 years or older) of the general community, in southern Brazil, aiming to contribute with epidemiological data on this disease in this age group.

METHOD

Study design and location

This is a cross-sectional, observational and retrospective study, developed with data obtained from exams registered in the system of a Molecular Diagnostics laboratory, which carried out the testing of suspected individuals for Covid-19, from July 2020 to May 2021, in public health services in

municipalities belonging to the Alto Jacuí and Alto da Serra do Botucaraí regions, which correspond to two of the twenty-eight regions that represent the Regional Development Councils (of Portuguese, Coredes), in the state of Rio Grande do Sul, Brazil²³.

The individuals treated in this laboratory were citizens of the following places: Boa Vista do Cadeado, Boa Vista do Incra, Colorado, Cruz Alta, Espumoso, Fontoura Xavier, Fortaleza dos Valos, Ibirubá, Lagoa dos Três Cantos, Mormaço, Não-Me-Toque, Quinze de Novembro, Saldanha Marinho, Santa Bárbara do Sul, Selbach, Tapera, Tupanciretã and Victor Graeff. The total number of residents of this group of cities is 192.967 people, and of this total, 29.395 are elderly people aged 60 years or older²⁴. Each of these municipalities had its own criteria for selecting individuals for testing, based on their epidemiological surveillance.

The diagnostic method adopted by the research laboratory was Charité²⁵ through detection of the virus E gene by the RT-qPCR technique, performed on nasopharyngeal and oropharyngeal swab samples. Samples were rejected when they didn't have an identification label, weren't accompanied by the exam request, had divergent information between the identification of the collection bottle and the exam request, had an incomplete number of swabs in the bottle or when the collection vials had some sort of breach.

For the exam result, the CT value (Cycle Threshold) of each sample was considered, positive samples being those with CT <35 for the E gene, inconclusive samples those with CT between 35 and 40 for the E gene and negative samples those without amplification, as previously validated by the laboratory.

Data collect

A retrospective survey was carried out of the data contained in the medical records of patients who underwent the test for Covid-19 using the RT-qPCR method, in the Molecular Diagnostics Laboratory of the University of Cruz Alta. The project was approved by the Research Ethics Committee, opinion No. 4,655,609. The secrecy and confidentiality of the data accessed in the laboratory records were ensured by the confidentiality agreement and the information collected was transcribed into a data collection instrument for exclusive use for the research, through a numerical system to catalog patient data, thus avoiding the use and identification of patients' names.

For the study, the following variables were collected: sex, age, body mass index (BMI), symptoms, comorbidities and RT-qPCR test results (dependent variable). Taking into account that the samples were collected and analyzed in a period when vaccination was not yet available or was only available for small groups, data regarding this variable were not collected.

Of the total of 3.872 exams performed in the research laboratory, only those referring to elderly patients aged 60 years or older were included in the present study, resulting in a final sample of 310 exams. Patients with inconclusive results were excluded from the research.

Statistical analysis

Data were compiled and analyzed using the SPSS® software (23.0 version, Chicago, IL Statistical Package for the Social Sciences). Percentage frequencies were calculated for all qualitative variables. Associations between these variables and positive results for Covid-19 were assessed using Pearson's chi-square or Fisher's exact tests, as appropriate. All estimates were bilateral with a pre-established level of significance for the alpha error of 5% ($p \leq 0.05$)

RESULTS

Of the 310 medical records of elderly patients evaluated, 75 (24.2%) had a positive RT-qPCR test for Sars-CoV-2. The age ranged from 60 to 99 years, with an average of 69 years (± 8.89) and there were a higher proportion of Covid-19 infection among elderly females (69.3%) compared to males (30.7%) ($p=0.017$) (Odds ratio =0.015). The BMI variable does not present statistically significant differences between the groups ($p=0.212$) (Table 1).

Table 1 – Characterization of elderly people negative and positive for Covid-19

Variables	Total		Covid-19 Negative		Covid-19 Positive		p* value
	No	%	No	%	No	%	
Sex							
Female	178	57.4	126	53.6	52	69.3	0.017
Male	132	42.6	109	46.4	23	30.7	
BMI**							
Underweight	4	1.6	3	1.5	1	1.8	0.212
Normal	66	25.6	51	25.2	15	26.8	
Overweight	85	32.9	61	30.2	24	42.9	
Obesity	103	39.9	87	43.1	16	28.6	

significant p values are highlighted

**The n does not correspond to the total of samples due to 52 requests not having this data filled in. BMI = body mass index

Regarding symptoms, 94% of the elderly had at least one clinical manifestation, as shown in table 2 that cough was reported by a total of 176 (56.2%) patients and that it was proportionally higher (67.1%) among the elderly with positive tests for Covid-19 ($p=0.028$) (Odds ratio =0.027). In addition, 148 (47.3%) of the elderly had fever $>37.8^{\circ}\text{C}$ in the laboratory records, and of these, the highest proportion (46.1%) was among those who tested positive for Sars-CoV-2 ($p=0.015$) (Odds ratio = 0.027) (Odds ratio =0.016).

Table 2 – Symptoms observed in negative and positive COVID-19 patients

Variables	Total		Covid-19 Negative		Covid-19 Positive		p* value
	No	%	No	%	No	%	
Cough	176	56.2	125	52.7	51	67.1	0.028
Coryza	148	47.3	111	46.8	37	48.7	0.779
Fever $>37^{\circ}\text{C}$	108	34.5	73	30.8	35	46.1	0.015
Dyspnea	45	14.4	34	14.3	11	14.5	0.978
Body ache	156	49.8	114	48.1	42	55.3	0.277
Sore throat	174	55.6	128	54.0	46	60.5	0.320
Diarrhea	72	23.0	56	23.6	16	21.1	0.642
Headache	132	42.2	96	40.5	36	47.4	0.292
Abdominal Pain	6	1.9	4	1.7	2	2.6	0.635

Otalgia	1	0.3	1	0.4	0	0.0	0.571
Inappetence	8	2.6	5	2.1	3	4.0	0.405
Nasal congestion	8	2.6	7	3.0	1	1.3	0.685
Ageusia	15	4.8	10	4.2	5	6.6	0.371
Anosmia	13	4.2	11	4.6	2	2.6	0.741
Shortness of breath	4	1.3	4	1.7	0	0.0	0.576
Fatigue	29	9.3	24	10.1	5	6.6	0.495
Vomit	28	8.9	22	9.3	6	7.9	0.820

significant *p* values are highlighted

Finally, Table 3 presents the main comorbidities presented by the individuals tested. It is observed that the most reported comorbidities were obesity (24.6%) and hypertension (13.4%). However, only heart diseases presented a significant *p* result when comparing the positive and negative Covid-19 for elderly groups (Odds ratio =0.028) (Table 3).

Table 3 – Comorbidities observed in negative and positive Covid-19 patients

Variables	Total		Covid-19 Negative		Covid-19 positive		<i>p</i> * value
	No	%	No	%	No	%	
Comorbidities	134	42.8	96	40.5	38	50.0	0.146
Diabetes	15	4.8	12	5.1	3	3.9	0.692
Hypertension	42	13.4	31	13.1	11	14.5	0.756
Heart disease	11	3.5	5	2.1	6	7.9	0.017
Autoimmune disease	1	0.3	0	0.0	1	1.3	0.243
Asthma	5	1.6	3	1.3	2	2.6	0.598
Rhinitis	2	0.6	2	0.8	0	0.0	0.422
Bronchitis	3	1.0	2	0.8	1	1.3	0.567
Obesity	77	24.6	54	22.8	23	30.3	0.188
Depression	1	0.3	1	0.4	0	0.0	0.571
Hypothyroidism	6	1.9	5	2.1	1	1.3	0.661
Immunosuppression	3	1.0	2	0.8	1	1.3	0.713

significant *p* values are highlighted

DISCUSSION

The present study demonstrates that 24.2% of elderly people presented a positive result in the RT-qPCR test for SARS-CoV-2, with an average age of 69 (±8.89) years and a predominance of positive results in females. Most participants had symptoms, especially cough and fever among the positive cases. Furthermore, the most frequent comorbidities were obesity and hypertension, however, heart disease was associated with positive cases.

In view of the recent pandemic scenario caused by the Sars-CoV-2 virus, responsible for the Covid-19 disease, understanding characteristics associated with the disease and its risk factors has

become fundamental for containing the pandemic and assisting the population. The elderly constitutes one of the main risk groups for morbidity and mortality from Covid-19 and our study identified that the disease was diagnosed in 24.2% of the elderly tested for investigation of this disease in public services in southern Brazil, which corroborates with the study published by Barbosa, Galvão, Souza, Gomes, Medeiros, Lima¹⁴ which demonstrated a proportion of Covid-19 cases in the state of Pará of 22.36%, similar to that found in our study.

There is evidence that the probability of infection by Sars-CoV-2 between men and women occurs in a similar way, however studies show a higher prevalence of positive cases in females, as observed in our study^{26,27}. This may be related to the fact that the entry of Sars-CoV-2 into cells occurs through the angiotensin-converting enzyme (ACE2), which is located on the X chromosome, and consequently, females have higher levels of this enzyme, which may influence the greater female susceptibility to infection^{27,28}. In addition, studies indicate that the elderly population makes great use of health services, and women seek doctor's appointments more due to several factors. Thus, another hypothesis to explain the greater occurrence of the disease in women would be due to the greater demand by the female public for health services and, consequently, the greater number of diagnoses of the disease^{29,30}.

We also identified a higher proportion of elderly people who had cough ($p=0.028$) and fever ($p=0.015$) among those with a positive test for the disease, corroborating other studies published in the literature^{1,31-33}, which cite fever and cough as initial and main symptoms of Sars-CoV-2 infection and serve as warning signs for the investigation of the disease as pointed out by the Pan American Health Organization (PAHO) together with the World Health Organization (WHO)³⁴.

Patients infected with SARS-CoV-2 often have a fever, with an elevation of body temperature above 37.8°C. This is due to the cytokine storm and the inflammatory reaction caused during infection, through the release of pro-inflammatory cytokines such as interleukin-1 (IL-1), interleukin-6 (IL-6) and tumor necrosis factor (TNF), which act as pyrogens causing the inflammatory response of fever³⁵⁻³⁶. In addition, cough is also a common symptom of Covid-19, arising from the ability of Sars-CoV-2 to infect peripheral sensory nerves in the vagus nerves, coughing mediators, leading to neuroinflammation and hypersensitivity mechanisms causing cough³⁷.

In addition, it is known that elderly patients have worse symptoms and lower survival rates due to Covid-19, this could be explained by the fact that with advancing age, the immune system remodels, becoming less responsive, which is crucial in fighting infections³⁸. Added to this, the elderly often has some type of comorbidity, which in turn ends up influencing the more severe progression of the disease, with hypertension, diabetes and heart disease being the most observed and described in the literature³⁹.

In our study, we observed a higher proportion of previous heart diseases among the elderly who were positive for Covid-19 than among those who were negative for the disease. This becomes relevant because studies confirm greater expression of the ACE2 gene in individuals with chronic diseases, which influences a greater chance of infection by SARS-CoV-2 and also a worse clinical course of the disease in patients with heart disease⁴⁰⁻⁴².

Our study had some limitations, among them, there was the difficulty in carrying out the return to the patients included in the study in order to verify the clinical outcome of those who had the disease. Furthermore, since data collection took place in a period when vaccination was not yet available, or was only available for some groups, data on immunization could not be included in the work. Furthermore, data collection (July 2020 to May 2021) comprised a period with various epidemiological scenarios of Covid-19 in Rio Grande do Sul and data analysis did not take into account whether the advancement of the age group would be related to more positive cases. Additionally, other

- ¹³ Guimarães AM, Reis LA, de Oliveira AS, Lopes AOS. Perfil epidemiológico de idosos acometidos pela Covid-19. *Saúde Coletiva* 2023;13(86).
- ¹⁴ Barbosa IR, Galvão MHR, Souza TAD, Gomes SM, Medeiros ADA, Lima KCD. Incidence and mortality from COVID-19 in the Brazilian elderly population and its relationship with contextual indicators: an ecological study. *Rev. Brazil geriatric Gerontol.* 2020;23(1):e200171.
- ¹⁵ Machado CJ, Pereira CCA, Viana BM, Oliveira GL, Melo DC, Garcia de Carvalho JFM, et al. Estimativas de impacto da COVID-19 na mortalidade de idosos institucionalizados no Brasil. *Ciência & Saúde Coletiva.* 2020;25(9):3.437-3.444. DOI: 10.1590/1413-81232020259.14552020
- ¹⁶ de Moraes EN, Viana LG, Resende LMH, Vasconcellos LS, Moura AS, Menezes A et al. COVID-19 nas instituições de longa permanência para idosos: estratégias de rastreamento laboratorial e prevenção da propagação da doença. *Ciência & Saúde Coletiva.* 2020;25(9):3.445-3.458. DOI: 10.1590/1413-81232020259.20382020
- ¹⁷ Araújo de Souza, de Araújo Nunes VM, do Nascimento ICS, Delmiro LAM, de Moaris MM, Nobre TTX et al. Vulnerabilidade e fatores de risco associados para Covid-19 em idosos institucionalizados. *Revista Eletrônica Acervo Saúde.* 2020;13(2)e5947. DOI: <https://doi.org/10.25248/reas.e5947.2021>
- ¹⁸ de Araújo PO, Freitas MYGS, Carvalho ESS, Peixoto TM, Servo MLS, Santana LS, et al. Institutionalized elderly: vulnerabilities and strategies to cope with Covid-19 in Brazil. *Educ. Enferm.* 2021;39(1):e07. DOI: <https://doi.org/10.17533/udea.iee.v39n1e07>
- ¹⁹ de Lacerda TTB, Assis MG, de Carvalho PF, Pereira LSM. Did managers of long-term care facilities neglect the functionality of older adults? A survey during the COVID-19 pandemic. *American Journal of Infection Control* 2023. DOI: <https://doi.org/10.1016/j.ajic.2023.05.001>
- ²⁰ Duarte MQ, Santo MAS, Lima CP, Giordani JP, Trentini CM. COVID-19 e os impactos na saúde mental: uma amostra do Rio Grande do Sul, Brasil. *Ciência & Saúde Coletiva.* 2020;25(9):3.401-3.411. DOI: 10.1590/1413-81232020259.16472020
- ²¹ Mafra TKA, Stobbe JC, Rabello RS, Lindemann IL, da Silva SG. A Síndrome Respiratória Aguda Grave na pessoa idosa no contexto da pandemia da covid-19 e seus fatores associados. *Rev. Bras. Geriatr. Gerontol.* 2023;26:e220158. DOI: <http://dx.doi.org/10.1590/1981-22562023026.220158.pt>
- ²² Novaes HPO, de Oliveira LMS, Gomes NP, Santos AA, Soub J, Duarte MB et al. Estratégias de enfrentamento à COVID-19 em instituições de longa permanência para idosos. *Rev Gaúcha Enferm.* 2023;44:e20210323. DOI: <https://doi.org/10.1590/1983-1447.2023.20210323.pt>
- ²³ Fórum dos Coredes. Conselhos Regionais de Desenvolvimento do Rio Grande do Sul. [Internet]. 2023. [access on 24 Sept. 2023]. Available at: <https://www.forumdoscoredes.org.br/>
- ²⁴ Instituto Brasileiro de Geografia e Estatística (IBGE). Censo Brasileiro de 2022. [Internet]. 2022. [access 20 Sept. 2023]. Available at: <https://cidades.ibge.gov.br/>
- ²⁵ Corman VM, Landt O, Kaiser M, Molenkamp R, Meijer A, Chu DKW, et al. detection of 2019 novel coronavirus (2019-nCoV) by real-time RT-PCR. *Euro Survey.* 2020;25(3):pii =2000045. DOI: <https://doi.org/10.2807/1560-7917>
- ²⁶ Fernández-de-Las-Peñas C, Martín-Guerrero JD, Pellicer-Valero ÓJ, Navarro-Pardo E, Gómez- Mayordomo V, Cuadrado ML, et al. Female Sex Is a Risk Factor Associated with long term Post-COVID Related-Symptoms but not with COVID-19 Symptoms: The LONG-COVID-EXP-CM Multicenter Study. *J Clin Med.* 2022 Jan. 14;11(2):413. DOI: 10.3390/jcm11020413
- ²⁷ Qian J, Zhao L, Ye RZ, Li XJ, Liu YL. Age dependent Gender Differences in COVID-19 in Mainland China: Comparative Study. *Clinic infect dis.* 2020 Dec. 3;71(9):2.488-2.494. DOI: 10.1093/cid /ciaa683
- ²⁸ Kelada M, Anto A, Dave K, Saleh SN. The Role of Sex in the Risk of mortality From COVID-19 Amongst adult Patients: The Systematic Review. *Cures.* 2020 Aug. 29;12(8):e10114. DOI: 10.7759/cureus.10114
- ²⁹ Pilger C, Menon MU, Mathias TADF. Use of health services by elderly people living in the community. *Journal of the School of Nursing at USP.* 2013;47:213-220. DOI: <https://doi.org/10.1590/S0080-62342013000100027>
- ³⁰ Turrini RNT, Cabrelli R, Iwanow Cianciarullo T, Silva Sousa C. Elderly in the family health unit: morbidity and use of health services. *Rev René [Internet].* 2014;15(1):89-98. [access May. 19, 2022]. Available at: <https://www.redalyc.org/pdf/3240/324030684012.pdf>
- ³¹ Cascella M, Ranik M, Aleem A, Dulebohn SC, Napoli RD. Features, Evaluation, and treatment of Coronavirus (COVID-19). *StatPearls [Internet].* 2022. [access May. 13 2022] Available at: <https://www.ncbi.nlm.nih.gov/books/NBK554776/>
- ³² Bhandari S, Shaktawat AS, Sharma R, Dube A, Kakkar S, Banerjee S, et al. A preliminary clinical-epidemiological portrayal of COVID-19 pandemic at a premier medical institution of North India. *Ann Thorac Med.* 2020 July-Sept;15(3):146-150. DOI: 10.4103/atm.ATM_182_20

- ³³ Rodriguez-Morales AJ, Cardona-Ospina JA, Gutiérrez-Ocampo E, Villamizar-Peña R, Holguin-Rivera Y, Escalera-Antezana JP, et al. Clinical, laboratory and imaging features of COVID-19: A systematic review and meta – analysis. *Travel Med Infect Dis.* 2020 Mar./Apr.;34:101623. DOI: 10.1016/j.tmaid.2020.101623
- ³⁴ Pan American Health Organization. Information sheet on COVID-19. [Internet]. 2022. [access May. 13, 2022]. Available at: <https://www.paho.org/pt/covid19>
- ³⁵ Balli S, Sharan S. Physiology, fever. In: StatPearls [Internet]. StatPearls Publication, 2020. [access May. 29, 2022]. Available at: <https://www.ncbi.nlm.nih.gov/books/NBK562334/>
- ³⁶ Conti P, Caraffa A, Tetè G, Gallenga CE, Ross R, Kritas SK, et al. mast cells activated by SARS-CoV- 2 release histamine which increases IL-1 levels causing cytokine storm and inflammatory reaction in COVID-19. *J Biol regulation homeost agents.* 2020 Sept./ Oct.;34(5):1.629-1.632. DOI: 10.23812/20-2EDIT
- ³⁷ Song WJ, Hui C, Hull JH, Birring SS, McGarvey L, Mazzone SB, et al. Confronting COVID-19-associated cough and the post-COVID syndrome: role of viral neurotropism, neuroinflammation, and neuroimmune responses. *Lancet Respir Med.* 2021;9(5):533-544. DOI: 10.1016/S2213-2600(21)00125-9
- ³⁸ Feng Y, Ling Y, Bai T, Xie Y, Huang J, Li J, et al. COVID-19 with Different Severities: A Multicenter study of Clinical Features. *Am J Breathe Crit Care Med.* 2020 June 1;201(11):1.380-1.388. DOI: 10.1164/rccm.202002-0445OC
- ³⁹ Guo T, Shen Q, Guo W, He W, Li J, Zhang Y, et al. Clinical Characteristics of Elderly patients with COVID-19 in Hunan Province, China: A Multicenter, Retrospective Study. *Gerontology.* 2020;66(5):467-475. DOI: 10.1159/000508734
- ⁴⁰ Moccia F, Gerbino A, Lionetti V, Miragoli M, Munaron LM, Pagliaro P, et al. COVID-19-associated cardiovascular morbidity in older adults: a position paper from the Italian Society of Cardiovascular Researches. *Geroscience.* 2020 Aug.;42(4):1021-1049. DOI: 10.1007/s11357-020-00198-w
- ⁴¹ Pathanhey G, Fadadu PP, Hospodar AR, Abbas AE. angiotensin-converting enzyme 2 and COVID-19: patients, comorbidities, and therapies. *Am J Physiol lung Cell Mol Physiol.* 2021 Mar. 1;320(3):L301-L330. DOI: 10.1152/ajplung.00259.2020
- ⁴² Gheblawi M, Wang K, Viveiros A, Nguyen Q, Zhong JC, Turner AJ, et al. Angiotensin-Converting Enzyme 2: SARS-CoV-2 Receptor and regulator of the Renin-Angiotensin System: Celebrating the 20th Anniversary of the Discovery of ACE2. *Circ Res.* 2020 May. 8;126(10):1.456-1.474. DOI: 10.1161/CIRCRESAHA.120.317015

Received: September 27, 2022

Accepted: November 21, 2023

Published: February 22, 2024

Authors' Contributions:

Luana Schayene Wagner: Conceptualization, Data curation, Formal analysis, Investigation, Visualization, Writing – original draft, Writing – review & editing

Gabrieli de Fátima Prudêncio dos Santos: Data curation, Writing – original draft, Investigation, Writing – review & editing

Juliana Lemes dos Santos: Data curation, Investigation, Validation, Writing – review & editing

Tatiana Mugnol: Data curation, Investigation, Validation, Writing – review & editing

Diego Pascoal Golle: Data curation, Investigation, Validation, Writing – review & editing

Josiane Woutheres Bortolotto: Data curation, Investigation, Validation, Writing – review & editing

Mariana Migliorini Parisi: Data curation, Investigation, Validation, Writing – review & editing

Janaina Coser: Conceptualization, Data curation, Formal analysis, Investigation, Project administration, Validation, Writing – original draft, Writing – review & editing

All authors have approved the final version of the text.

Financing: Does not have financing.

Conflict of interest: There are no conflicts of interest.

Corresponding Author:

Luana Schayene Wagner
Universidade de Cruz Alta – Unicruz
Rodovia Municipal Jacob Della Mea, s/n km 5,6- Parada Benito
Cruz Alta/RS, Brasil. CEP 98020-290
E-mail: luana.wagner@sou.unicruz.edu.br

Editor: Dr. Matias Nunes Frizzo

Editor-in-chief: Dra. Adriane Cristina Bernat Kolankiewicz

This is an open access article distributed under the terms of the Creative Commons license.

