

ORIGINAL ARTICLE

## Association of C-Reactive Protein with Cardiovascular Risk Factors, Lung Function and Quality of Life in Young Adults

Natalia Silva Bueno<sup>1</sup>; Pedro Henrique de Almeida Silva<sup>2</sup>; Ana Julia Ribeiro Gomes<sup>3</sup>  
Carolina Silva Carvalho<sup>4</sup>; Eron Matheus Leite Moreira<sup>5</sup>  
Sarah Rhaquel Rodrigues Oliveira<sup>6</sup>; Viviane Soares<sup>7</sup>

Highlight

Increased inflammatory response is associated with obesity;

Increased inflammatory response affects lung function;

Psychological, functional and vitality quality of life were higher in young adults with moderate cardiovascular risk.

ABSTRACT

**Objective:** To associate ultrasensitive C-reactive protein (US-CRP) with cardiovascular risk factors, lung function and quality of life. **Methods:** Body mass index (BMI), waist circumference (WC), haemodynamic parameters, blood sampling for US-CRP, spirometry (peak expiratory flow – PEF; forced expiratory volume in one second – FEV<sub>1</sub>) and individualised application of the Short Form-36 (SF-36) health-related quality of life (HRQoL) questionnaire. **Results:** Of the 100 young adults studied, BMI and WC were higher in those at high risk. PEF and FEV<sub>1</sub>/FVC ratio were higher in low-risk young adults. Mental health, functional capacity and vitality (VT) were higher in young people at lower cardiovascular risk. Of the overweight/obese young adults, 87.5% were at moderate and high risk (p=0.044), and of those with WC above the predicted level, 86.7% were at moderate and high risk (p=0.002). Among young adults with HRQoL above P(50), 43.5%, 50%, and 45.7% were at low cardiovascular risk for functional capacity, general health, and vitality, respectively. US-CRP was positively correlated with BMI (p=0.044), whereas PEF (p=0.046) and FEV<sub>1</sub>/FVC ratio (p=0.002) were inversely correlated. **Conclusions:** US-CRP was associated with BMI and WC, lung function and HRQoL, especially with respect to physical aspects. Obesity control should be emphasized, and regular physical activity and proper nutrition may help, including reducing the chronic inflammatory response.

**Keywords:** inflammation; obesity; abdominal obesity; pulmonary function test; quality of life

<sup>1</sup> Universidade Evangélica de Goiás, Curso de Medicina. Anápolis/GO, Brasil. <https://orcid.org/0000-0002-8195-074X>

<sup>2</sup> Universidade Evangélica de Goiás, Programa de Pós-Graduação em Movimento Humano e Reabilitação. Anápolis/GO, Brasil. <http://orcid.org/0000-0002-1832-6776>

<sup>3</sup> Universidade Evangélica de Goiás, Curso de Medicina. Anápolis/GO, Brasil. <http://lattes.cnpq.br/0380398757985637>

<sup>4</sup> Universidade Evangélica de Goiás, Curso de Medicina. Anápolis/GO, Brasil. <https://orcid.org/0000-0001-5961-9623>

<sup>5</sup> Universidade Evangélica de Goiás, Curso de Medicina. Anápolis/GO, Brasil. <https://orcid.org/0009-0008-8057-9812>

<sup>6</sup> Universidade Evangélica de Goiás, Curso de Medicina. Anápolis/GO, Brasil. <https://orcid.org/0000-0002-5332-8566>

<sup>7</sup> Universidade Evangélica de Goiás. Programa de Pós-Graduação em Movimento Humano e Reabilitação. Anápolis/GO, Brasil. <https://orcid.org/0000-0002-1570-6626>

## INTRODUCTION

Currently, the increasing prevalence of overweight has become a global public health challenge.<sup>1</sup> Obesity is a chronic disease in which the accumulation of adipose tissue leads to consequences such as an increase in cardiovascular and metabolic disease, disability and mortality, and is associated with low levels of physical activity.<sup>3,4</sup> The most commonly used markers of obesity are body mass index (BMI) and waist circumference (WC), the latter being used as a marker of abdominal obesity.<sup>4</sup>

Obesity is an important cause of cardiovascular disease (CVD), and several factors have been studied to elucidate the relationship between the two conditions.<sup>5</sup> Among them, C-reactive protein (CRP) has been shown to be an influential predictor of CVD, as adipose tissue can influence the secretion of pro-inflammatory cytokines, and when these enter the systemic circulation and reach the liver, they stimulate the release of CRP.<sup>6</sup> Thus, a low chronic inflammatory response is a common aspect of obese individuals.<sup>7</sup>

In addition, the accumulation of fat in the body causes changes in respiratory physiology, affecting various parameters of lung function, such as reduced diaphragmatic mobility and costal movement.<sup>8</sup> The association between obesity and reduced lung function is related to low levels of physical activity and poorer quality of life (QoL) in obese individuals.<sup>4,9</sup> There is also an association between elevated CRP, obesity and the presence of restrictive ventilatory disorders.<sup>10</sup> The overlap of these conditions leads to impairment in aspects related to QoL, both physical (activities of daily living and vitality) and mental health.<sup>3</sup>

In summary, it is suggested that elevated CRP is a current risk factor, is associated with obesity, leads to a low chronic inflammatory response, and that this condition may affect lung function and have consequences for HRQoL, especially aspects related to activities of daily living and vitality. The literature on the relationship between CRP and lung function and HRQoL is still in its infancy. The literature that has been found is based on adult samples, whereas the present study is based on young people. If CRP levels are high in young people, there is likely to be obesity and, as a result, impaired lung function and quality of life. If these conditions persist into adulthood, there is a risk of developing metabolic and cardiovascular disease and restrictive ventilatory disorders. The aim of this study was to investigate the association of ultrasensitive C-reactive protein (US-CRP) with cardiovascular risk factors, lung function and quality of life.

## METHODS

This is an analytical, cross-sectional study carried out with young people enrolled in a higher education course in the field of health at a university. The study was carried out between August 2018 and February 2019, with 180 students enrolled from the first to the tenth semester. Regularly enrolled young people aged 18 and over were included, while those with chronic co-morbidities (heart disease, chronic lung disease) and cognitive impairment registered with the course secretary were excluded. The sample was recruited by convenience and power was calculated using G\*Power 3.1 software in post-hoc mode, considering the analysis performed (correlation of continuous variables), an effect size ( $\rho_{H1}$ ) of 0.3, a significance level of 5%, and a power of 92%.

The study was approved by the Human Research Ethics Committee of the Evangelical University of Goiás (CEP-Unievangélica), following the guidelines of Resolution 466/2012 of the National Health Council (CNS), according to opinion number 2.989.854/2018. All participants signed an informed consent form (ICF). The young adults were approached by the researchers during class, with the permission and schedule of the teacher in the class at the time. The stages and assessments were explained by reading the ICF orally and then giving it to the young people to read again and sign. This

was followed by an explanation of the blood collection procedures, which were scheduled according to the participant's availability. On the same day, the QoL questionnaire was administered, and anthropometric and haemodynamic measurements were taken. The lung function test was performed on a different day because of the need for preparation: no coffee on the day of the test, no physical activity in the previous 24 hours, and no smoking in the previous 2 hours. The tests were carried out at the university, specifically in the multidisciplinary laboratory of the physiotherapy course.

Sociodemographic data were collected using an identification form with information on sex, alcohol and smoking history, physical activity and current medication.

BMI (kg/m<sup>2</sup>) was calculated by dividing body mass (kg) (Filizola digital scale, model 2096 PP, São Paulo, Brazil) by height (m) squared (Sanny stadiometer, São Paulo, Brazil). The cut-off points for eutrophic and overweight/obese subjects were  $\leq 24.99$  kg/m<sup>2</sup> and  $\geq 25$  kg/m<sup>2</sup>, respectively.<sup>11</sup> WC was measured using an inextensible anthropometric tape (Teklife, model TL200, São Paulo, Brazil) positioned at the midpoint between the upper iliac crest and the last costal arch at the end of expiration at rest. The reference values to characterise WC as within the predicted range were less than 94 cm for men and 80 cm for women.<sup>12</sup>

Systemic blood pressure was measured with the young adults sitting for five minutes at rest. Measurements were taken one minute apart, and the first measurement was discarded.<sup>13</sup> A semi-automatic device (OMRON, model HEM 705CP, Kyoto, Japan) was used to measure systolic blood pressure (SBP) and diastolic blood pressure (DBP). The reference values for normal SBP and DBP were less than 120 x 80 mmHg.

A blood sample was taken for US-CRP measurement (5 mL taken to analyse 1 mL of serum). The method of analysis was immunoturbidimetric and classified as low ( $< 1$  mg/dL), intermediate (1 - 3 mg/dL) and high ( $> 3$  mg/dL) cardiovascular risk.<sup>14</sup>

Spirometry was performed using a portable device (brand Mir, model MiniSpir, Rome, Italy) with specific disposable filters and mouthpieces. The participant performed the test in a seated position with the head in a neutral position and attached to a nose clip. Acceptability and reproducibility criteria were based on the American Thoracic Society/European Respiratory Society standard.<sup>15</sup> Manoeuvres performed with 1) maximal inspiration before the test, 2) satisfactory start of expiration, 3) evidence of maximal effort, 4) test duration of at least 6 seconds, 5) plateau in the last second and 6) absence of cough, leak, obstruction, Valsalva manoeuvre and glottis closure were accepted. The parameters analysed were peak expiratory flow (PEF), forced expiratory volume in one second (FEV<sub>1</sub>), forced vital capacity (FVC) and FEV<sub>1</sub>/FVC ratio. For reproducibility, two manoeuvres were considered if the difference in PEF was  $< 0.5$  L and FEV<sub>1</sub> if the difference between manoeuvres was  $< 0.15$  L. The results were interpreted according to the values predicted for the Brazilian population.<sup>16</sup>

HRQoL was assessed using the Short Form-36 (SF-36) questionnaire.<sup>17</sup> The questionnaire has a multi-item scale that includes: functional capacity (10 items), limitations due to physical health problems (4 items), limitations due to mental/emotional health problems (3 items), social functioning (2 items), emotional well-being (5 items), pain (2 items), vitality (4 items), and perception of general health (5 items). Scores range from 0 to 100, with lower scores corresponding to poorer HRQoL and higher scores reflecting better QoL.<sup>17</sup>

The results are presented as mean, standard deviation, frequencies, percentages, and variation ( $\Delta$ ) between means. Normality of the data was tested using the Kolmogorov-Smirnov test. Comparisons between groups were made using the one-way Anova test with post hoc for variables with a normal distribution and the Kruskal-Wallis test with Dunn's post hoc for variables with an asymmetric distribution. The chi-squared test was used to test the association between categorical variables, and the SF-36 domains were classified according to percentiles (P50). The correlation between continuous

variables was assessed using Pearson's coefficient (normal distribution) and Spearman's coefficient (asymmetric distribution). The p-value used was  $< 0.05$  and the data were analyzed using the Statistical Package for Social Science (SPSS, IBM, version 23.0, Armonk, NY) software.

## RESULTS

The characterisation of the sample is described in Table 1. Most of the young adults were female (82%), without a history of alcoholism (51%) or smoking (79%) and without regular physical activity (62%). The most common medication used was contraceptives (17%).

Table 1 – Characterization of the sample of young adults who took part in the study (n = 100)

Variables	Mean (SD)
Age (years)	20,90±2,37
Mass (kg)	62,04±11,16
Height (cm)	166,26±08,04
	<b>n (%)</b>
<b>Sex</b>	
Women	82 (82)
Male	18 (18)
<b>Ethylism</b>	49 (49)
<b>Smoking</b>	21 (21)
<b>Physical activity</b>	37 (37)
<b>Medicines in use</b>	
Anxiolytics/antidepressants	10 (10)
Contraceptives	17 (17)
Analgesia/anti-inflammatory	3 (3)
Vitamins and minerals	4 (4)
Neurological	4 (4)
Intestinal disorders	5 (5)
Others	5 (5)

Source: Prepared by the authors (2023).

With regard to anthropometric data, young adults at high cardiovascular risk had higher BMI (low risk:  $\Delta = +3.78 \text{ kg/m}^2$ ,  $p < 0.001$ ; moderate risk:  $\Delta = +2.77 \text{ kg/m}^2$ ,  $p = 0.011$ ) and WC (low risk:  $\Delta = +6.90 \text{ cm}$ ,  $p = 0.001$ ; moderate risk:  $\Delta = +5.53 \text{ cm}$ ,  $p = 0.019$ ) (Table 2).

About lung function, young adults at moderate cardiovascular risk had a higher PEF compared to those at low ( $\Delta = +1.74 \text{ L/s}$ ,  $p < 0.001$ ) and high risk ( $\Delta = +1.10 \text{ L/s}$ ,  $p = 0.013$ ). In addition, young adults classified as low risk according to CRP had a higher  $FEV_1/FVC$  ratio compared to those at moderate ( $\Delta = +0.06$ ,  $p = 0.021$ ) and high risk ( $\Delta = +0.04$ ,  $p = 0.037$ ).

In terms of perceived quality of life, young people at moderate cardiovascular risk had a higher mental health score than those at high cardiovascular risk ( $\Delta = +13.85$ ,  $p = 0.037$ ). In terms of functional capacity, young people at moderate risk had high scores compared with young people at low ( $\Delta = +7.36$ ,  $p = 0.041$ ) and high ( $\Delta = +8.65$ ,  $p = 0.007$ ) risk. Finally, participants at low cardiovascular risk had higher scores in the vitality domain compared to those at moderate risk ( $\Delta = +5.93$ ,  $p = 0.007$ ).

Table 2 – Comparison of anthropometric measurements, hemodynamic parameters, pulmonary function and quality of life according to CRP (n = 100)

Variables	CRP			p*
	< 1 mg/dL (n = 48)	1 – 3 mg/dL (n = 26)	> 3 mg/dL (n = 26)	
<b>Anthropometric measurements</b>				
BMI (kg/m <sup>2</sup> )	21.18±2.60	22.19±3.13	24,96±4,55	<0,001
WC (cm)	69,75±6,70	71,12±7,72	76,65±7,83	0,001
<b>Hemodynamic parameters</b>				
SBP (mmHg)	106,23±11,56	105,73±14,29	107,04±10,61	0,925
DBP (mmHg)	69,83±7,09	71,50±9,05	73,96±7,59	0,097
<b>Pulmonary function</b>				
PEF (L/s)	6.16±1.97	4.42±1.51	5.52±1.51	0.001
FEV <sub>1</sub> (L)	3.07±0.71	2.93±0.68	3.08±0.70	0.435
FVC (L)	3.25±0.77	3.30±0.72	3.38±0.78	0.774
FEV <sub>1</sub> /FVC ratio	0.95±0.09	0.89±0.13	0.91±0.10	0.027
<b>Domains of the SF-36</b>				
Mental health	56.08±20.77	59.08±17.32	45.23±20.70	0.030
Functional capacity	80.52±18.63	87.88±17.95	79.23±12.94	0.021
Physical limitations	64.58±36.41	61.54±33.34	67.31±33.74	0.778
Emotional limitation	42.36±41.11	44.87±37.64	43.59±41.92	0.930
Social aspects	62.80±25.56	44.87±37.64	43.59±41.92	0.349
Pain	59.10±24.72	63.31±23.73	60.31±15.24	0.655
General state of health	53.25±13.87	57.23±15.54	48.58±13.49	0.071
Vitality	45.42±22.12	51.35±18.03	36.54±17.76	0.026

Legend: BMI: body mass index; WC: waist circumference; SBP: systolic blood pressure; DBP: diastolic blood pressure; PEF: peak expiratory flow; FEV<sub>1</sub>: forced expiratory volume in the first second; FVC: forced vital capacity. \*Data for p < 0.05.

Source: Prepared by the authors (2023).

There was a significant association between CRP and BMI (p < 0.001) and WC (p = 0.002). Of the overweight/obese young adults, 87.5% were at moderate and high risk, and of those with WC above the predicted level, 86.7% had CRP ≥ 1 mg/dL (moderate and high risk). Among young people with HRQoL above P(50), 43.5%, 50% and 45.7% were at low cardiovascular risk for functional capacity, general health and vitality, respectively.

Table 3 – Association of anthropometric variables, hemodynamic parameters, pulmonary function and quality of life according to CRP (n=100)

Variáveis	CRP			p*
	< 1 mg/dL (n = 48) n (%)	1 – 3 mg/dL (n = 26) n (%)	> 3 mg/dL (n = 26) n (%)	
<b>BMI (kg/m<sup>2</sup>)</b>				
Eutrophic	45 (59.2)	20 (26.3)	11 (14.5)	< 0.001
Overweight/Obesity	03 (12.5)	06 (25.0)	15 (62.5)	
<b>WC (cm)</b>				
On schedule	46 (54.1)	22 (25.9)	17 (20.0)	0.002
Above expectations	02 (13.3)	04 (26.7)	09 (60.0)	

<b>SBP (mmHg)</b>				
Normotensive	48 (49.0)	24 (24.5)	26 (26.5)	0.064
Hypertensive	00 (00.0)	02 (100.00)	00 (0.00)	
<b>DBP (mmHg)</b>				
Normotensive	47 (50.0)	23 (24.5)	24 (25.5)	0.226
Hypertensive	01 (16.7)	03 (50.0)	02 (33.3)	
<b>FEV<sub>1</sub> (L)</b>				
On schedule	12 (41.4)	08 (27.6)	09 (31.0)	0.667
Below expectations	36 (50.7)	18 (25.4)	17 (23.9)	
<b>FVC (L)</b>				
On schedule	15 (40.5)	12 (32.4)	10 (27.0)	0.441
Below expectations	33 (52.4)	14 (22.2)	16 (25.4)	
<b>FEV<sub>1</sub>/FVC ratio</b>				
On schedule	46 (50.5)	21 (23.1)	24 (26.4)	0.115
Below expectations	02 (22.2)	05 (55.6)	02 (22.2)	
<b>Mental health</b>				
> P (50)	21 (47.7)	15 (34.1)	08 (18.2)	0.148
< P (50)	27 (48.2)	11 (19.6)	18 (32.1)	
<b>Functional capacity</b>				
> P (50)	20 (43.5)	18 (39.1)	08 (17.4)	0.015
< P (50)	28 (51.9)	08 (14.8)	18 (33.3)	
<b>Physical limitations</b>				
> P (50)	19 (54.3)	07 (20.0)	09 (25.7)	0.551
< P (50)	29 (44.6)	19 (29.2)	17 (26.2)	
<b>Emotional limitation</b>				
> P (50)	18 (43.9)	10 (24.4)	13 (31.7)	0.554
< P (50)	30 (50.8)	16 (27.1)	13 (22.0)	
<b>Social aspects</b>				
> P (50)	23 (48.9)	14 (29.8)	10 (21.3)	0.531
< P (50)	25 (47.2)	12 (22.6)	16 (30.2)	
<b>Pain</b>				
> P (50)	18 (45.0)	12 (30.0)	10 (25.0)	0.755
< P (50)	30 (50.0)	14 (23.3)	16 (26.7)	
<b>General state of health</b>				
> P (50)	25 (50.0)	17 (34.0)	08 (16.0)	0.041
< P (50)	23 (46.0)	09 (18.0)	18 (36.0)	
<b>Vitality</b>				
> P (50)	21 (45.7)	17 (37.0)	08 (17.4)	0.040
< P (50)	27 (50.0)	09 (16.7)	18 (33.3)	

Legend: BMI: body mass index; WC: waist circumference; SBP: systolic blood pressure; DBP: diastolic blood pressure; PEF: peak expiratory flow; FEV<sub>1</sub>: forced expiratory volume in the first second; FVC: forced vital capacity; P: percentile. \*Data for p < 0.05.

Source: Prepared by the authors (2023).

CRP had a positive correlation with BMI ( $p = 0.044$ ), while PEF ( $p = 0.046$ ) and the FEV<sub>1</sub>/FVC ratio ( $p = 0.002$ ) showed an inversely proportional correlation.

Table 4 – Correlation between CRP and anthropometric and pulmonary function measurements (n=100)

Variáveis	CRP (mg/dL)	
	r	p*
BMI (kg/m <sup>2</sup> )	0.202	0.044
PEF (L/s)	-0.200	0.046
FEV <sub>1</sub> /FVC ratio	-0.302	0.002

Legend: BMI: body mass index; PEF: peak expiratory flow; FEV<sub>1</sub>: forced expiratory volume in the first second; FVC: forced vital capacity. \*Data for p < 0.05.

Source: Prepared by the authors (2023).

## DISCUSSION

The main findings of the study were that obesity, BMI and WC were higher in young adults with CRP > 3 mg/dL, while PEF and FEV<sub>1</sub>/FVC ratio were higher in young adults with low cardiovascular risk. In terms of quality of life, scores were higher for mental health, functional capacity and vitality, with the first two domains being higher in young people with CRP between 1 and 3 mg/dL, and the third domain being higher in young people with low-risk CRP. Most of the young adults who were obese were at moderate and high cardiovascular risk. CRP was positively correlated with BMI, while PEF and FEV<sub>1</sub>/FVC ratio were negatively correlated.

Young adults with CRP > 3 mg/dL had higher body mass index, BMI and WC, and total CRP levels were directly correlated with BMI. These results are supported by the literature, as in young people CRP is closely associated with anthropometric indicators, BMI and WC.<sup>18,19</sup> This is justified because obesity is signaled by an inflammatory state in which there is an increase in adipose tissue together with a decrease in adiponectin. Associated with this process, adipose tissue is taken over by inflammatory cytokines, leading to local inflammation and consequently to systemic inflammation, defined by a marked increase in inflammatory markers such as CRP.<sup>18, 20</sup>

In terms of lung function, PEF and FEV<sub>1</sub>/FVC ratio were higher and inversely related in students with low-risk CRP. Consistent with these findings, a greater decline in FEV<sub>1</sub> has been reported in individuals with higher baseline CRP in young adults in their 20s.<sup>10,21</sup> In addition, it has been shown that lung function suffers with increased inflammation.<sup>22</sup> However, in the study by Ahmadi-Abhari et al.<sup>23</sup> a significant association between CRP and lung function was found only in older individuals, as CRP is a marker of an age-related pro-inflammatory process, but may have high concentrations in young people depending on their health-related factors. According to this theory, chronic inflammation increases telomere shortening, which in turn leads to senescence of the alveolar and endothelial cells of the lung. In addition, endothelial dysfunction induced by systemic inflammation can lead to pulmonary vascular infiltration and lung tissue damage.<sup>23</sup>

Scores were higher in the domains of mental health, functional capacity, and vitality, with the first two domains being higher in young people with CRP between 1 and 3 mg/dL, whereas the third domain was higher in low-risk participants. Taking into account the questions of the QoL questionnaire (SF-36), it can be concluded that a decline in general health already has an impact on QoL. Consistent with these findings, the study by Cho et al.<sup>24</sup> states that the higher the plasma concentration of CRP, the greater the fatigue, independent of risk factors such as BMI, depressive symptoms, sleep quality, pain and physical activity. Fatigued individuals may be less physically active, and low physical activity may lead to an increase in CRP levels. This is because physical and psychological stressors activate the peripheral immune system and trigger an inflammatory response with the release of pro-inflammatory cytokines and acute phase proteins.<sup>25</sup> It is worth noting that the level of physical activity was not the subject of this study but should be considered in future studies.

The following limitations should be noted. Firstly, it was not possible to establish a cause-and-effect relationship because of the cross-sectional nature of the study. In addition, there was low adherence among young people, but the sampling power, based on the type of data analysis carried out, was sufficient to generalize the results to the population studied. Secondly, many young adults were unable to reach the six-second value for the forced expiratory maneuver, but this was minimized by the ATS guidance using estimates. In terms of strengths, the study is innovative in relating CRP to HRQoL, as it was conducted in young people who theoretically would not have elevated blood CRP due to the direct relationship with age (young adults rarely have elevated CRP), and the study of the relationship between CRP and lung function parameters is important, as airway obstruction leads to greater inflammation and consequently higher CRP concentrations.

## CONCLUSION

It can therefore be concluded that there is a directly proportional relationship between CRP and BMI, whereas the relationship between CRP and lung function is inversely proportional. The HRQoL assessment showed higher scores in mental health and functional capacity in young people with CRP between 1 and 3 mg/dL, and in vitality in young people with low cardiovascular risk. However, further studies are needed to clarify the mechanical and physiological effects of CRP on lung function. In this sense, it is important to control obesity to reduce the inflammatory response, and to this end, regular physical activity and an adequate diet can help to reduce the chronic inflammatory response.

## REFERENCES

- <sup>1</sup> Chooi YC, Ding C, and Magkos F. The epidemiology of obesity. *Metabolism*. 2019;92:6-10. Available at: <https://www.sciencedirect.com/science/article/abs/pii/S002604951830194X>
- <sup>2</sup> Wang Y, Beydoun MA, Min J, Xue H, Kaminsky LA, and Cheskin LJ. Has the prevalence of overweight, obesity and central obesity levelled off in the United States? Trends, patterns, disparities, and future projections for the obesity epidemic. *Int J Epidemiol*. 2020;49(3):810-823. DOI: 10.1093/ije/dyz273
- <sup>3</sup> Dalle Grave R, Soave F, Ruocco A, Dametti L, and Calugi S. Quality of Life and Physical Performance in Patients with Obesity: A Network Analysis. *Nutrients*. 2020;12(3). DOI: 10.3390/nu12030602
- <sup>4</sup> Svartengren M, et al. The impact of body mass index, central obesity and physical activity on lung function: results of the EpiHealth study. *ERJ Open Res*. 2020;6(4). DOI: 10.1183/23120541.00214-2020
- <sup>5</sup> Powell-Wiley TM, et al. Obesity and Cardiovascular Disease: A Scientific Statement From the American Heart Association. *Circulation*. 2021;143(21):p. e984-e1010. DOI: 10.1161/cir.0000000000000973
- <sup>6</sup> Pavela G, Kim Y-i, and Salvy S-J. Additive effects of obesity and loneliness on C-reactive protein. *PloS one*. 2018;13(11):e0206092. DOI: 10.1371/journal.pone.0206092
- <sup>7</sup> Saltiel AR and Olefsky JM. Inflammatory mechanisms linking obesity and metabolic disease. *J Clin Invest*. 2017;127(1):1-4. DOI: 10.1172/jci92035
- <sup>8</sup> Brock JM, Billeter A, Müller-Stich BP, and Herth F. Obesity and the Lung: What We Know Today. *Respiration*. 2020;99(10):856-866. DOI: 10.1159/000509735
- <sup>9</sup> Ishikawa C, Barbieri MA, Bettiol H, Bazo G, Ferraro AA, and Vianna EO. Comparison of body composition parameters in the study of the association between body composition and pulmonary function. *BMC Pulm Med*. 2021;21(1):178. DOI: 10.1186/s12890-021-01543-1
- <sup>10</sup> Jung DH, Shim JY, Ahn HY, Lee HR, Lee JH, and Lee YJ. Relationship of body composition and C-reactive protein with pulmonary function. *Respir Med*. 2010;104(8):1.197-1.203. DOI: 10.1016/j.rmed.2010.02.014
- <sup>11</sup> World Health Organization. Diet, nutrition, and the prevention of chronic diseases: report of a joint WHO/FAO expert consultation. World Health Organization; 2003; Vol. 916.
- <sup>12</sup> World Health Organization. Waist Circumference and Waist – Hip Ratio. Report of a WHO Expert Consultation. Geneva; 8-11 Dec. 2008. 2008:8-11.
- <sup>13</sup> Barroso WKS, et al. Diretrizes Brasileiras de Hipertensão Arterial–2020. *Arquivos brasileiros de cardiologia*. 2021;116:516-658. DOI: 10.36660/abc.20201238



- <sup>14</sup> Ridker PM. Cardiology Patient Page. C-reactive protein: a simple test to help predict risk of heart attack and stroke. *Circulation*. 2003;108(12):e81-5. DOI: 10.1161/01.cir.0000093381.57779.67
- <sup>15</sup> Graham BL, et al. Standardization of spirometry 2019 update. An official American thoracic Society and European respiratory Society technical statement. *American journal of respiratory and critical care medicine*. 2019;200(8):e70-e88. DOI: 10.1164/rccm.201908-1590ST
- <sup>16</sup> Neder JA, Andreoni S, Castelo-Filho A, and Nery LE. Reference values for lung function tests. I. Static volumes. *Braz J Med Biol Res*. 1999;32(6):703-17. DOI: 10.1590/s0100-879x1999000600006
- <sup>17</sup> Ciconelli RM, Ferraz MB, Santos W, Meinao IM, and Quaresma MR. Tradução para a língua portuguesa e validação do questionário genérico de avaliação de qualidade de vida SF-36 (Brasil SF-36). *Revista Brasileira de Reumatologia*. 1999;39(3):143-150. Available at: <https://pesquisa.bvsalud.org/portal/resource/pt/lil-296502>
- <sup>18</sup> Cercato C and Fonseca FA. Cardiovascular risk and obesity. *Diabetology & Metabolic Syndrome*. 2019;11(1):74. DOI: <https://doi.org/10.1186/s13098-019-0468-0>
- <sup>19</sup> Menezes CA, Costa GLOB, Barreto RF, and Oliveira VS. Proteína C reativa importante biomarcador de risco cardiometabólico na obesidade infanto-juvenil. *Saúde Coletiva (Barueri)*. 2021;11(65):5.882-5.895. DOI: <https://doi.org/10.36489/saudecoletiva.2021v11i65p5882-5895>
- <sup>20</sup> Landecho MF, Tuero C, Valentí V, Bilbao I, de la Higuera M, and Frühbeck G. Relevance of Leptin and Other Adipokines in Obesity-Associated Cardiovascular Risk. *Nutrients*. 2019;11(11):2.664. DOI: 10.3390/nu11112664
- <sup>21</sup> Rasmussen F, et al. High-sensitive C-reactive protein is associated with reduced lung function in young adults. *Eur Respir J*. 2009;33(2):382-388. DOI: 10.1183/09031936.00040708
- <sup>22</sup> Nybo M, Hansen HS, Siersted HC, and Rasmussen F. No relationship between lung function and high-sensitive C-reactive protein in adolescence. *The Clinical Respiratory Journal*. 2010;4(4):230-236. DOI: 10.1111/j.1752-699X.2009.00181.x
- <sup>23</sup> Ahmadi-Abhari S, Kaptoge S, Luben RN, Wareham NJ, and Khaw KT. Longitudinal association of C-reactive protein and lung function over 13 years: The EPIC-Norfolk study. *Am J Epidemiol*. 2014;179(1):48-56. DOI: 10.1093/aje/kwt208
- <sup>24</sup> Cho HJ, Seeman TE, Bower JE, Kiefe CI, and Irwin MR. Prospective association between C-reactive protein and fatigue in the coronary artery risk development in young adults study. *Biol Psychiatry*. 2009;66(9):871-878. DOI: 10.1016/j.biopsych.2009.06.008
- <sup>25</sup> Black PH. Stress and the inflammatory response: a review of neurogenic inflammation. *Brain Behav Immun*. 2002;16(6):622-653. DOI: 10.1016/s0889-1591(02)00021-1

Submitted: January 16, 2023

Accepted: November 30, 2023

Published: May 14, 2024

### Contribuições dos autores

Natalia Silva Bueno: conceptualization, investigation, methodology, project administration, writing original drafting.

Pedro Henrique de Almeida Silva: data validation, design of data presentation, writing – proofreading and editing, translation.

Ana Julia Ribeiro Gomes: conceptualization, investigation, methodology, project administration, writing original drafting.

Carolina Silva Carvalho: conceptualization, investigation, methodology, project administration, writing original drafting.

Eron Matheus Leite Moreira: conceptualization, investigation, methodology, project administration, writing original drafting.

Sarah Rhaquel Rodrigues Oliveira: conceptualization, investigation, methodology, project administration, writing original drafting.

Viviane Soares: conceptualization, investigation, methodology, project administration, resources, supervision, visualization, writing – review & editing.

---

**All authors approved the final version of the text.**

**Conflicts of interest:** There are no conflicts of interest.

**Financing:** Does not have financing

**Corresponding author**

Viviane Soares

Curso de Medicina

Programa de Pós-Graduação em Movimento Humano e Reabilitação

Universidade Evangélica de Goiás – Unievangélica

Av. Universitária, s/n - Cidade Universitária, Anápolis/GO, Brasil.

ftviviane@gmail.com

**Associate Editor:** Eliane Roseli Winkelmann (Ph.D)

**Editor-in-Chief:** Adriane Cristina Bernat Kolankiewicz (Ph.D)

This is an open-access article distributed under the terms of the Creative Commons License

