

**LIFESTYLE HABITS, PHYSICAL DISABILITY, AND THE RELATIONSHIP  
WITH NON-SPECIFIC CHRONIC LOW BACK PAIN**

Mikael Gleidison de Melo Araújo,<sup>1</sup> Etiene Oliveira da Silva Fittipaldi,<sup>2</sup>  
Albanita Gomes da Costa de Ceballos,<sup>3</sup> Washington José dos Santos<sup>4</sup>

Highlights: (1) The intensity of low back pain is associated with greater physical disability. (2) Smoking increases the risk of low back pain and physical disability. (3) Lack of leisure activities is linked to greater low back pain and physical disability.

PRE-PROOF

(as accepted)

This is a preliminary, unedited version of a manuscript that has been accepted for publication in the journal Context & Health. As a service to our readers, we are making this early version of the manuscript available as accepted. The article will still be reviewed, formatted and approved by the authors before it is published in its final form.

<http://dx.doi.org/10.21527/2176-7114.2024.48.14452>

How to cite:

Araújo MG de M, Fittipaldi EO da S, de Ceballos AG da C, dos Santos WJ. Lifestyle habits, physical disability, and the relationship with chronic non-specific low back pain. Rev. Contexto & Saúde, 2024;24(48): e14452

---

<sup>1</sup> Prefeitura da Cidade do Recife. Recife/PE, Brasil. <https://orcid.org/0009-0009-8977-5921>

<sup>2</sup> Universidade Federal de Pernambuco, Recife/PE, Brasil. <https://orcid.org/0000-0002-1524-6930>

<sup>3</sup> Universidade Federal de Pernambuco, Recife/PE, Brasil. <https://orcid.org/0000-0002-8658-9981>

<sup>4</sup> Empresa Brasileira de Serviços Hospitalares, Hospital das Clínicas da Universidade Federal de Pernambuco, Recife/PE, Brasil. <https://orcid.org/0000-0002-2291-8563>

## LIFESTYLE HABITS, PHYSICAL DISABILITY, AND THE RELATIONSHIP WITH NON-SPECIFIC CHRONIC LOW BACK PAIN

### ABSTRACT

**Objective:** This study examined the relationship between lifestyle habits, physical disability, and non-specific chronic low back pain. **Methods:** This cross-sectional, exploratory, and analytical study included a convenience sample of 50 individuals aged  $\geq 18$  years of both sexes. Participants were assessed on socioeconomic, demographic, and lifestyle factors, as well as the level of physical disability, using the Roland-Morris questionnaire and pain intensity using the visual analog scale. **Results:** The participants had an average age of 56.9 (SD = 8.87), ranging from 34 to 72 years. Among them, 90% (n = 45) were female. Pain perception was divided between moderate (30%, n = 15) and severe (70%, n = 35). There was a statistically significant association between pain and physical disability (prevalence ratio [PR] 9.86, confidence interval [CI] = 1.46–66.47). Individuals who did not engage in leisure activities also reported a higher subjective feeling of low back pain (PR 0.60, CI = 0.37–0.95) and physical disability (PR 0.39, CI = 0.18–0.88). Additionally, there was a statistically significant association between smoking and LBP (PR 1.60, CI = 1.25–1.99), as well as between physical disability and low back pain (PR 1.88, CI = 1.13–3.11). No significant associations were found between physical exercise, alcohol consumption, and diet with levels of pain and physical disability. **Conclusion:** A strong relationship was found between pain and physical disability, as well as smoking habits and interference with leisure activities.

**Keywords:** Low back pain. Chronic pain. Physical disability. Lifestyle habits.

### INTRODUCTION

Low back pain (LBP) is a musculoskeletal disorder that significantly affects the quality of life, independence, and social participation of individuals<sup>1</sup>. This condition also imposes a high burden on public services, with epidemic levels in the general population<sup>2</sup>. The prevalence of LBP is higher in industrialized countries, affecting approximately 37% of the global adult population<sup>3</sup>. Data from the Brazilian National Health Survey conducted in 2019<sup>4</sup> revealed that around 21.6% of the Brazilian population experiences some form of chronic back problem, most commonly in the lumbar region. Moreover, it is estimated that 80% of adults will experience LBP at some point in their lives<sup>5</sup>.

## **LIFESTYLE HABITS, PHYSICAL DISABILITY, AND THE RELATIONSHIP WITH NON-SPECIFIC CHRONIC LOW BACK PAIN**

According to Hartvigsen et al.<sup>3</sup>, it is not possible to identify a specific nociceptive cause for chronic low back pain (CLBP). There is strong evidence that sociodemographic factors, lifestyle choices, educational level, psychosocial or work-related factors, and their interactions influence the onset and persistence of this condition. Evidence indicates that the physical disability caused by CLBP is not solely attributed to biological factors. Despite its high prevalence rate, approximately 90–95% of CLBP cases have no identifiable etiology and are multifactorial<sup>6</sup>.

Understanding the factors contributing to CLBP and their impact on individuals is crucial for healthcare professionals to effectively manage this issue. It is important to consider not only the factors directly associated with CLBP but also the subjective aspects associated with this condition. This knowledge enables the design of appropriate multidimensional interventions. Therefore, this study aims to examine the relationship between lifestyle habits, physical disability, and CLBP.

### **METHODOLOGY**

This exploratory cross-sectional study was conducted in three Family Health Units (FHUs) in the Santo Amaro neighborhood in the central area of Recife (Pernambuco State, northeastern Brazil). The study took place between April 5 and December 20, 2022. Participants were included based on convenience, without a sample calculation. Participants were recruited from professionals of the family health strategy, informed about the study's objective, and tasked with identifying individuals in the area who met the criteria. These individuals were then invited to the unit to participate in the study at a designated day and time. Additionally, participants were attracted through health education sessions conducted by the researcher in the waiting rooms of the FHUs.

The inclusion criteria were as follows: being 18 years or older, experiencing CLBP for at least three months, and being registered at one of the FHUs involved in the study. Exclusion criteria included cognitive and/or neurological impairments, previous neuromusculoskeletal dysfunctions (e.g., trauma sequelae, amputations, and deformities), and pregnancy.

Participants received both verbal and written explanations about the study and were asked to sign the informed consent form in duplicate; one copy was retained by the participant, while the other remained with the researcher. Data collection occurred in a private area,

## **LIFESTYLE HABITS, PHYSICAL DISABILITY, AND THE RELATIONSHIP WITH NON-SPECIFIC CHRONIC LOW BACK PAIN**

specifically the reception room of the FHU, to maintain confidentiality and respect the participants' privacy.

The research project was approved by the Research Ethics Committee (protocol no. 5.384.447). It adhered to the ethical guidelines recommended by Resolution no. 466/2012 of the National Health Council of the Brazilian Ministry of Health pertaining to research involving human subjects.

### **Procedures**

Initially, participants completed a questionnaire prepared by the researchers. The questionnaire included questions about socioeconomic and demographic factors, such as race (white, black, brown), sex, age, education level (illiterate, elementary school, high school), occupation, marital status (with or without a partner), and family income (up to one Brazilian minimum wage or more than one minimum wage). Additionally, the questionnaire included questions about lifestyle habits, such as regular physical exercise, alcohol use, smoking, and consumption of processed foods, fruits, and vegetables.

Next, the researchers administered the Roland-Morris questionnaire, which measures the impact of LBP on activities of daily living (ADLs) and instrumental activities of daily living (IADLs). The questionnaire has been validated for Brazilian Portuguese and consists of 24 questions. It is a quick and easy assessment that takes about five minutes to complete. Scores are calculated by adding up the responses, ranging from zero (no disability) to 24 (severe disability). Scores above 14 indicate physical disability<sup>7</sup>.

Finally, the subjective sensation of pain was assessed using the Visual Analog Scale, which quantifies pain intensity based on an individual's perception. The scale consists of different levels of pain intensity, and participants are asked to verbally indicate their pain level using corresponding phrases<sup>8</sup>. Pereira et al.<sup>9</sup> suggested a pain score classification, with 0 representing no pain, mild pain ranging from 1 to 2, moderate pain ranging from 3 to 7, and severe pain ranging from 8 to 10.

### **Statistical Analysis**

The instruments were converted into an online form using Google Forms. For data analysis, numerical variables were reported as means and standard deviations, while categorical

**LIFESTYLE HABITS, PHYSICAL DISABILITY, AND THE RELATIONSHIP  
WITH NON-SPECIFIC CHRONIC LOW BACK PAIN**

variables were presented as frequencies and percentages. Since this study was cross-sectional, the prevalence ratio (PR) was calculated to determine the association between pain sensation and degree of disability, as well as the relationship between lifestyle habits and pain sensation and disability. A 95% confidence interval (CI) was considered. The data was analyzed using the Statistical Package for the Social Sciences (SPSS) software, version 22.0.

## RESULTS

A total of 50 individuals were assessed, of whom 90% (n = 45) were female with an average age of 56.9 (SD = 8.87) years, ranging from 34 to 72 years. Among the respondents, 62% (n = 31) identified themselves as brown and 50% (n = 25) reported having partners. In terms of education, 52% (n = 26) had not completed elementary school. Regarding family income, 76% (n = 38) received less than one minimum monthly wage. In terms of occupation, 56% (n = 28) identified themselves as “housewives.” None of the individuals reported feeling mild pain, with the perception of pain being divided between moderate (30%) and severe (70%) (n = 35) (Table 1).

Table 1 - Sociodemographic and pain intensity variables of individuals with chronic low back pain from three family health units in Recife (Pernambuco, northeastern Brazil), 2022.

Variable	Moderate pain	Severe pain	n (%)
<b>Sex</b>			
Female	14	31	45 (90)
Male	1	4	5 (10)
<b>Race/color</b>			
White	4	7	10 (22)
Black	3	5	8 (16)
Brown	8	23	31 (62)
<b>Marital status</b>			
With a partner	8	17	25 (50)
Without a partner	7	18	25 (50)
<b>Education</b>			
Illiterate	0	3	3 (6)
Elementary education	9	17	26 (52)
High school education	6	15	21 (42)
<b>Occupation</b>			
Health worker	0	4	4 (8)
Retired	2	1	3 (6)
Businessman	0	1	1 (2)

**LIFESTYLE HABITS, PHYSICAL DISABILITY, AND THE RELATIONSHIP  
WITH NON-SPECIFIC CHRONIC LOW BACK PAIN**

Seamstress	1	0	1 (2)
Housewife	8	20	28 (56)
Student	0	1	1 (2)
Mechanic	1	0	1 (2)
Driver	0	1	1 (2)
Stonemason	0	1	1 (2)
Service provider	2	5	10 (14)
Security guard	1	0	1 (2)
Unemployed	0	1	1 (2)
<b>Household income</b>			
≤1 minimum wage	10	28	38 (76)
>1 minimum wage	5	7	12 (24)

According to the classification of the disability instrument, 48% (n = 24) of the individuals fell into the disability group, while 52% (n = 26) did not have any disability. The prevalence ratio between severe pain and disability was 9.86 (CI = 1.46-66.47), as shown in Table 2. The mean score for pain sensation was 8.06 (SD = 1.43; min = 5, max = 10). In the questionnaire, the mean score was 12.92 (SD = 6.09; min = 2, max = 24).

Table 2 - Prevalence ratio between pain and physical disability in individuals with chronic low back pain from three family health units in Recife (Pernambuco, northeastern Brazil), 2022.

Sensation of pain	Physical disability – n (%)		PR (CI = 95%)
	Yes	No	
Severe	23 (46)	12 (24)	9.86 (1.46–66.47)
Moderate	1 (2)	14 (28)	

PR: Prevalence ratio; CI: confidence interval.

After all 50 participants answered the questions on lifestyle habits, the results showed that 62% (n = 31) reported not consuming alcohol, 94% did not exercise (n = 47), 18% were smokers (n = 9), 40% participated in leisure activities (n = 20), and 72% ate processed foods (n = 36). As for the consumption of fruit and vegetables, 70% confirmed doing so (n = 35). Table 3 presents the relationship between LBP and lifestyle habits. The findings showed a statistically significant association between smoking and pain intensity as well as leisure activities, with a PR of 1.60 (CI = 1.46–66.47) for the former and PR of 0.60 (CI = 0.37–0.95) for the latter.

**LIFESTYLE HABITS, PHYSICAL DISABILITY, AND THE RELATIONSHIP  
WITH NON-SPECIFIC CHRONIC LOW BACK PAIN**

Table 3 - Prevalence ratio between lifestyle habits and pain in patients with chronic low back pain from three family health units in Recife (Pernambuco, northeastern Brazil), 2022.

Lifestyle habits	Sensation of pain – n (%)		PR (CI = 95%)
	Moderate	Severe	
Alcohol consumption	15 (30)	4 (8)	1.22 (0.86–1.73)
Smoking	9 (18)	0 (0)	1.60 (1.25–1.99)
Physical exercise	2 (4)	1 (2)	0.95 (0.42–2.16)
Leisure activities	10 (20)	10 (20)	0.60 (0.37–0.95)
Industrialized food consumption	26 (52)	10 (20)	1.12 (0.72–1.74)
Fruit and vegetable consumption	12 (24)	3 (6)	1.21 (0.85–1.72)

PR: Prevalence ratio; CI: confidence interval.

Table 4 lists the association between physical disability and lifestyle habits. It was possible to observe statistically significant associations between physical disability and smoking and leisure time activity, respectively PR = 1.88 (CI = 1.13–3.11) and PR = 0.39 (CI = 0.18–0.88).

Table 4 - Prevalence ratio between lifestyle habits and physical disability of patients with chronic low back pain from three family health units in Recife (Pernambuco, northeastern Brazil), 2022.

Lifestyle habits	Physical disability – n (%)		PR (CI = 95%)
	Yes	No	
Alcohol consumption	10 (20)	9 (18)	1.16 (0.65–2.07)
Smoking	7 (14)	2 (4)	1.88 (1.13–3.11)
Physical exercise	0 (0)	3 (6)	—
Leisure activities	5 (10)	15 (30)	0.39 (0.18–0.88)
Industrialized food consumption	16 (32)	20 (40)	0.79 (0.43–1.39)
Fruit and vegetable consumption	7 (14)	8 (16)	0.96 (0.51–1.82)

PR: Prevalence ratio; CI: confidence interval

## LIFESTYLE HABITS, PHYSICAL DISABILITY, AND THE RELATIONSHIP WITH NON-SPECIFIC CHRONIC LOW BACK PAIN

### DISCUSSION

Most participants in this study were females with low levels of education and income up to one minimum wage. There was a higher prevalence of severe low back pain and disability among these participants. Smokers had a higher prevalence of disability, while there was an inverse relationship between leisure activity and disability. In the literature, socioeconomic factors are commonly cited as risk factors for the generation and maintenance of chronic pain<sup>10,11</sup>. According to a study by Malta et al.<sup>12</sup>, based on 2019 data<sup>4</sup>, back pain is highly prevalent and is associated with demographic factors and socioeconomic conditions.

Although the mean age of the participants in this study was over 50, there were women who were not yet menopausal, and the different phases of the menstrual cycle can also affect the perception of low back pain. A study by Carvalho and Souza<sup>13</sup> showed a higher prevalence of chronic pain in women compared to men, which also affects pain intensity and ADLs. Souza, Häfele, and Siqueira<sup>14</sup> observed that chronic pain perception was higher among women, with the lumbar spine being one of the most affected regions.

This study found a high prevalence of individuals with low levels of education. According to Romero et al.<sup>15</sup>, adults with lower levels of education (no education or incomplete elementary education) have a higher prevalence of chronic back problems compared to those with higher levels of education. The same finding was reported in a study by Rocha, Alfieri, and Silva<sup>16</sup>, as most participants with chronic pain had a lower level of education.

Furthermore, this study found that individuals with CLBP had a higher prevalence of physical disabilities and limitations in ADLs and IADLs. Chronic pain significantly interferes with everyday activities and significantly impacts the quality of life<sup>17</sup>. The years lived with disability caused by low back pain have increased by over 50% since 1990, especially in low- and middle-income countries<sup>18</sup>. Trelha et al.<sup>19</sup> observed that chronic pain mainly affects the ability to get up (72%), perform household tasks (54%), and walk (47%), suggesting that severe pain interferes with an individual's level of functionality. Lemos et al.<sup>20</sup> also indicated a relationship between ADLs and pain intensity.

Notably, 94% of the participants in this study did not engage in exercise, and despite there being no apparent relationship between physical exercise, pain, and disability, research has demonstrated an important association between a sedentary lifestyle and CLBP<sup>21</sup>. Evidence



## **LIFESTYLE HABITS, PHYSICAL DISABILITY, AND THE RELATIONSHIP WITH NON-SPECIFIC CHRONIC LOW BACK PAIN**

shows that the level of physical activity influences pain intensity, indicating that sedentary or insufficiently active individuals report higher pain intensity compared to those who are active and very active<sup>22</sup>. More than half of the individuals reported frequent consumption of processed foods. Although no relationship was found, the literature suggests that industrialized and high-fat foods play a role in pain generation by inducing the formation of inflammatory substances in the body<sup>12,23</sup>.

Another important finding in this study was the lower prevalence of severe pain and physical disability among individuals who reported engaging in leisure activities. This suggests that there may be a relationship between leisure activities and pain. Pereira et al.<sup>9</sup> conducted a study that showed that individuals with severe CLBP tended to experience more physical disability, which impaired their social interaction and, consequently, their ability to engage in leisure activities. Trelha et al.<sup>19</sup> found that pain mainly interfered with sleep (61.28%), mood (54.25%), and leisure activities (45.05%). Scudds and Ostbye<sup>24</sup> conducted a study on a population of elderly Canadians, revealing that pain interfered with mood in approximately 54% of the participants, with leisure activities in 44%, and with sleep in 40%.

Moreover, our findings showed a higher prevalence of severe pain among smokers compared to non-smokers. Malta et al.<sup>12</sup>, using official government data<sup>4</sup>, assessed the prevalence of back pain among smokers, ex-smokers, and non-smokers and noted that back pain was present in the majority of smokers; Caputo<sup>25</sup> also observed an association between back pain and smoking. CLBP affects all age groups and is generally associated with sedentary occupations, smoking, obesity, and low socioeconomic status<sup>18</sup>. Hence, these findings suggest a possible relationship between the intensity of CLBP and disability in smokers, emphasizing the need for further research on this topic.

### **LIMITATIONS OF THE STUDY**

The study has inherent limitations in its design. Age may have influenced the study results, as older individuals are more prone to musculoskeletal dysfunctions. In this study, the participants' ages varied. Additionally, there is a possibility of overestimating the prevalence rates since CLBP was self-reported, and the answers were based on the participants' perceptions of their lifestyle habits and physical disability. Another limitation is the lack of sample calculation, as the survey was conducted for convenience rather than representing the

**LIFESTYLE HABITS, PHYSICAL DISABILITY, AND THE RELATIONSHIP  
WITH NON-SPECIFIC CHRONIC LOW BACK PAIN**

entire population. This convenience sampling may have resulted in the exclusion of mild cases of CLBP.

## **CONCLUSION**

It is evident from this and other findings in the literature that chronic low back pain cannot be approached and understood solely through the lens of nociception as it is also related to socioeconomic factors, demographic factors, and lifestyle habits. In this study, a strong correlation was observed between pain and physical disability, as well as smoking habits and interference with leisure activities. Thus, primary healthcare professionals have tremendous potential to promote healthy habits through multidisciplinary and interdisciplinary approaches to develop effective strategies to reduce the comorbidities and disabilities caused by chronic low back pain.

## **REFERENCES**

- <sup>1</sup> Stefane T, Santos AM, Marinovic A, Hortense P. Chronic low back pain: pain intensity, disability and quality of life. *Acta Paul Enferm.* 2013;26(1):14-20.
- <sup>2</sup> Carregaro RL, da Silva EN, van Tulder M. Direct healthcare costs of spinal disorders in Brazil. *Int J Public Health.* 2019;64(6):965-974.
- <sup>3</sup> Hartvigsen J, Hancock MJ, Kongsted A, Louw Q, Ferreira ML, Genevay S. et al. What low back pain is and why we need to pay attention. *Lancet.* 2018;391(10137):2356-2367.
- <sup>4</sup> INSTITUTO BRASILEIRO DE GEOGRAFIA E ESTATÍSTICA (IBGE). Pesquisa Nacional de Saúde: 2019: Percepção do estado de saúde, estilos de vida, doenças crônicas e saúde bucal: Brasil e grandes regiões. IBGE, Coordenação de Trabalho e Rendimento; 2019.
- <sup>5</sup> Refshauge KM, Maher CG. Low back pain investigations and prognosis: a review. *Br J Sports Med.* 2008;40(6):494-498.
- <sup>6</sup> Krismer M, van Tulder M. Low back pain (non-specific). *Best Pract Res Clin Rheumatol.* 2007;21:77-91.
- <sup>7</sup> Nusbaum L, Natour J, Ferraz MB, Goldenberg J. Translation, adaptation and validation of the Roland-Morris questionnaire-Brazil Roland- Morris. *Brazilian Journal of Medical and Biological Research.* 2001;34(2):203-210.
- <sup>8</sup> Batalha LMC. Avaliação da dor. Coimbra: ESEnfC; 2016 (Manual de estudo versão 1).

**LIFESTYLE HABITS, PHYSICAL DISABILITY, AND THE RELATIONSHIP  
WITH NON-SPECIFIC CHRONIC LOW BACK PAIN**

<sup>9</sup> Pereira LV, Vasconcelos PP de, Souza LAF, Pereira G de A, Nakatani AYK, Bachion MM. Prevalence and intensity of chronic pain and self- perceived health among elderly people: a population-based study. *Rev Lat Am Enfermagem*. 2014;22(4):662-669.

<sup>10</sup> Kiadaliri A, Merlo J, Englund M. Complex sociodemographic inequalities in consultations for low back pain: lessons from multilevel intersectional analysis. *Pain*. 2021;162(4):1135-1143.

<sup>11</sup> Malta DC, Oliveira MM de, Andrade SSC de A, Caiaffa WT, Souza M de FM de, Bernal RTI. Factors associated with chronic back pain in adults in Brazil. *Rev Saude Publica*. 2017;51 Suppl 1:9s.

<sup>12</sup> Malta DC, Bernal RTI, Ribeiro EG, Ferreira E de MR, Pinto RZ, Pereira CA. Chronic back pain among Brazilian adults: data from the 2019 National Health Survey. *Rev. bras. epidemiol*. 2022;25:e220032.

<sup>13</sup> Carvalho AD, Souza EP. O Idoso e as Dores Crônicas: como Viver com Elas. *Rev. Mult. Psic (on line)*. 2017;11(38):689-700.

<sup>14</sup> Souza DF da S de, Häfele V, Siqueira FV. Dor crônica e nível de atividade física em usuários das unidades básicas de saúde. *Rev Bras Ativ Fís Saúde*. 2019;24:e0085.

<sup>15</sup> Romero DE, Santana D, Borges P, Marques A, Castanheira D, Rodrigues JM, Sabbadini L. Prevalência, fatores associados e limitações relacionados ao problema crônico de coluna entre adultos e idosos no Brasil. *Cad. Saúde Pública*. 2018;34(2):e00012817.

<sup>16</sup> Rocha ADX, Alfieri FM, Silva NCOV. Prevalence of chronic pain and associated factors in a small town in southern Brazil. *BrJP. São Paulo*, 2021 jul-sep;4(3):225-231.

<sup>17</sup> Donzeli MA, Magalhaes LF, Oliveira GVA, Dias AA, Gasparini ALC, Bertonecello D. Nível de incapacidade e qualidade de vida em mulheres com dor lombar crônica. *REFACS (online)*. 2020;8(2):261-266.

<sup>18</sup> Clark S, Horton R. Low back pain: a major global challenge. *Lancet*. 2018;391(10137):2302.

<sup>19</sup> Trelha CS, Panazzolo D, Dellaroza MSG, Cabrera MAS, Souza R, Pisconti F, Taho YM. Capacidade funcional de idosos com dor crônica residentes na comunidade. *Geriatrics & Gerontologia*. 2008;2(2):59-64.

<sup>20</sup> Lemos BO, Cunha AMR, Cesarino CB, Martins MRI. The impact of chronic pain on functionality and quality of life of the elderly. *BrJP*. 2019;2(3):237-241.

<sup>21</sup> Pereira Júnior AA, Benvenuti A. Dor Lombar em Mulheres Sedentárias e Praticantes de Musculação. *Cinergis*. 2016;18(1):54-58.

<sup>22</sup> Ferretti F, Silva MRD, Pegoraro F, Baldo JE, Sá CAD. Chronic pain in the elderly, associated factors and relation with the level and volume of physical activity. *BrJP*. 2019;2(1):3-7.

**LIFESTYLE HABITS, PHYSICAL DISABILITY, AND THE RELATIONSHIP  
WITH NON-SPECIFIC CHRONIC LOW BACK PAIN**

<sup>23</sup> Boyd JT, LoCoco PM, Furr AR et al. Elevated dietary  $\omega$ -6 polyunsaturated fatty acids induce reversible peripheral nerve dysfunction that exacerbates comorbid pain conditions. *Nat Metab.* 2021;3:762-773.

<sup>24</sup> Scudds RJ, Ostbye T. Pain and pain-related interference with function in older Canadians: the canadian study of health and aging. *Disabil Rehabil.* 2001;23(15):654-664.

<sup>25</sup> Caputo EL, Souza DF da S de, Häfele V, Siqueira FCV. Back pain prevalence and associated factors in Brazilian Unified Health System users. *BrJP.* 2022;5(2):137-142.

Submitted: May 10, 2023

Accepted: November 29, 2023

Published: April 22, 2024

**Authors' contributions:**

Mikael Gleidison de Melo Araújo – Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Writing – original draft, Writing – review & editing

Etiene Oliveira da Silva Fittipaldi - Writing – original draft, Writing – review & editing

Albanita Gomes da Costa de Ceballos - Writing – original draft, Writing – review & editing

Washington José dos Santos - Conceptualization, Data curation, Formal analysis, Methodology, Supervision, Writing – original draft, Writing – review & editing

All the authors approved the final version of the text.

Conflict of interest: There is no conflict of interest.

Financing: Does not have financing

**LIFESTYLE HABITS, PHYSICAL DISABILITY, AND THE RELATIONSHIP  
WITH NON-SPECIFIC CHRONIC LOW BACK PAIN**

**Corresponding author:**

Washington José dos Santos

Hospital das Clínicas da Universidade Federal de Pernambuco

Av. Prof. Moraes Rego, 1235 - Cidade Universitária, Recife/PE, Brasil. CEP50670-901

E-mail: [washingtonfisio@gmail.com](mailto:washingtonfisio@gmail.com)

Editor: Dr. Eliane Roseli Winkelmann

Editor-in-chief: Dr. Adriane Cristina Bernat Kolankiewicz

This is an open-access article distributed under the terms of the Creative Commons license.



PRE-PROOF