

**BROMATOLOGICAL ANALYSIS AND EVALUATION OF ANTIOXIDANT  
PROPERTIES OF PASTEURIZED CUPUAÇU (*Theobroma grandiflorum*) JUICE  
FERMENTED BY *Lacticaseibacillus rhamnosus* ATCC 9595**

Daniel dos Santos Guimarães<sup>1</sup>; Camila Caetano da Silva<sup>2</sup>  
Kátia Sayuri Aragão Aguiar<sup>3</sup>; Alexia Figueiredo Ferreira<sup>4</sup>; Marcos Andrade Silva<sup>5</sup>  
Luís Cláudio Nascimento da Silva<sup>6</sup>; Adrielle Zagnignan<sup>7</sup>

**Highlights:** (1) Fermentation by *L. rhamnosus* ATCC 9595 increased the antioxidant capacity of cupuaçu juice. (2) The juice met microbiological standards, with *L. rhamnosus* remaining stable for 28 days. (2) Fermented juice maintained antioxidant activity during storage for 28 days.

PRE-PROOF

(as accepted)

This is a preliminary, unedited version of a manuscript that has been accepted for publication in Revista Contexto & Saúde. As a service to our readers, we are making this initial version of the manuscript available, as accepted. The article will still be reviewed, formatted, and approved by the authors before being published in its final form.

<http://dx.doi.org/10.21527/2176-7114.2024.48.15050>

How to cite:

Guimarães D dos S, da Silva CC, Aguiar KSA, Ferreira AF, Silva MA. et al. Bromatological analysis and evaluation of antioxidant properties of pasteurized cupuaçu (*Theobroma grandiflorum*) juice fermented by *Lacticaseibacillus rhamnosus* ATCC 9595. Rev. Contexto & Saúde, 2024;24(48): e15050

---

<sup>1</sup> Universidade Ceuma. São Luís/MA, Brasil. <https://orcid.org/0009-0007-6699-596X>

<sup>2</sup> Universidade Ceuma. São Luís/MA, Brasil. <https://orcid.org/0009-0003-2830-7821>

<sup>3</sup> Universidade Ceuma. São Luís/MA, Brasil. <https://orcid.org/0009-0000-1550-2132>

<sup>4</sup> Universidade Ceuma. São Luís/MA, Brasil. <https://orcid.org/0009-0000-1550-2132>

<sup>5</sup> Universidade Ceuma. São Luís/MA, Brasil. <https://orcid.org/0000-0001-6558-2863>

<sup>6</sup> Universidade Ceuma. São Luís/MA, Brasil. <https://orcid.org/0000-0002-4206-0904>

<sup>7</sup> Universidade Ceuma. São Luís/MA, Brasil. <https://orcid.org/0000-0001-9865-2223>

**BROMATOLOGICAL ANALYSIS AND EVALUATION OF ANTIOXIDANT PROPERTIES OF  
PASTEURIZED CUPUAÇU (*Theobroma grandiflorum*) JUICE FERMENTED BY  
*Lactiseibacillus rhamnosus* ATCC 9595**

## **ABSTRACT**

This study conducted a bromatological analysis and evaluated the antioxidant properties of pasteurized cupuaçu (*Theobroma grandiflorum*) juice fermented by *Lactiseibacillus rhamnosus* ATCC 9595. The juices were pasteurized (80 °C/10 minutes) for subsequent fermentation with *L. rhamnosus* ATCC 9595 (inoculum of 10<sup>8</sup> CFU/mL). After 48 hours, the viability of *L. rhamnosus*, production of organic acids, and resistance to lysozyme and bile salts were analyzed. The samples were refrigerated for 28 days for bromatological analyses. The juices were extracted with ethyl acetate to evaluate antioxidant activity, phenolic compounds, and flavonoids. *L. rhamnosus* ATCC 9595 grew in pasteurized cupuaçu juice (~9 Log CFU/mL) and remained stable over 28 days ( $p > 0.05$ ). All samples met the microbiological standards established by Brazilian guidelines for juices. Similarly, no significant changes were detected in the levels of ashes, moisture, or Brix degrees during the analyzed periods ( $p > 0.05$ ). *L. rhamnosus* ATCC 9595 cultivated in the juice or MRS medium resisted the action of lysozyme (100 mg/L) and bile salts (0.5% and 1.0%). The antioxidant capacity of cupuaçu juice significantly increased after fermentation by *L. rhamnosus* ATCC 9595, as did the levels of phenolic compounds. The fermented juice samples showed similar IC<sub>50</sub> values during storage, while the non-fermented extract showed variation from the 21st day onwards. The results indicate that fermentation by *L. rhamnosus* ATCC 9595 is an efficient strategy to enhance the antioxidant characteristics of cupuaçu juice.

**Keywords:** Fermentation, Fruit Juice, Probiotics, Functional Food, Free Radicals.

## **1. INTRODUCTION**

Free radicals and reactive oxygen species (ROS) play important roles in the etiology of degenerative pathologies (Parkinson's disease, Alzheimer's disease), cancer, diabetes, and inflammatory disorders<sup>1</sup>. In this sense, organisms have developed various strategies to control these agents, constituting the antioxidant defense system formed by enzymes (such as superoxide dismutase and catalase) and proteins (glutathione and thioredoxin)<sup>2</sup>. However, in many situations, this system cannot cope with the overproduction of reactive species, leading to a state called oxidative stress related to the clinical manifestations described above<sup>3,4</sup>.

**BROMATOLOGICAL ANALYSIS AND EVALUATION OF ANTIOXIDANT PROPERTIES OF  
PASTEURIZED CUPUAÇU (*Theobroma grandiflorum*) JUICE FERMENTED BY  
*Lacticaseibacillus rhamnosus* ATCC 9595**

An alternative way to delay the damage caused by free radicals is by using foods with antioxidant properties that act through different mechanisms <sup>1,5</sup>. Fruits are considered excellent sources of bioactive compounds (such as proteins, vitamins, soluble carbohydrates, flavonoids, and other phenolic compounds) <sup>6</sup>. For example, cupuaçu juice and pulp (*Theobroma grandiflorum*) possess compounds with antioxidant, anti-inflammatory, and hypoglycemic properties <sup>7-9</sup>.

Recently, the development of a potentially probiotic cupuaçu juice was reported based on the incorporation of *Lacticaseibacillus rhamnosus* ATCC 9595 (= *Lactobacillus rhamnosus*) <sup>9</sup>. This strain was selected due to its antimicrobial and immunomodulatory properties <sup>10,11</sup>. Fermentation of cupuaçu juice with *L. rhamnosus* ATCC 9595 caused alterations in the metabolite profile of cupuaçu juice due to biotransformation and an increased concentration of some bioactive compounds <sup>9</sup>.

Additionally, fermentation by *L. rhamnosus* ATCC 9595 increased the anti-inflammatory properties observed in murine models of lipopolysaccharide-induced endotoxemia <sup>9</sup>. Fermentation with *L. rhamnosus* ATCC 9595 was also employed to improve the functional characteristics of bacuri juice. The fermented bacuri (*Platonia insignis*) juice showed superior anti-infectious effects compared to the non-fermented juice, prolonging the survival of *Tenebrio molitor* larvae infected with an enteroaggregative *Escherichia coli* strain <sup>12</sup>.

Given the biotechnological potential of *L. rhamnosus* ATCC 9595 for the production of potentially probiotic beverages, this study conducted a bromatological analysis and evaluated the antioxidant properties of pasteurized cupuaçu juice fermented by *L. rhamnosus* ATCC 9595 stored for 28 days.

## **2. MATERIALS AND METHODS**

### **2.1. Preparation of pasteurized juices and fermentation with *Lacticaseibacillus rhamnosus* ATCC 9595**

Cupuaçu fruits were obtained in São Luís (Maranhão, Brazil). The fruit pulp was manually removed and stored at -20 °C until preparation. In each experiment, samples (30 g) of pulp were dissolved in 250 mL of distilled water (120 mg/mL), and the pH was adjusted to 6.0. Then, each juice was subjected to pasteurization (80 °C for 10 min). The juice was

**BROMATOLOGICAL ANALYSIS AND EVALUATION OF ANTIOXIDANT PROPERTIES OF  
PASTEURIZED CUPUAÇU (*Theobroma grandiflorum*) JUICE FERMENTED BY  
*Lacticaseibacillus rhamnosus* ATCC 9595**

transferred to a container with ice until it reached room temperature, and then 1 mL of *L. rhamnosus* ATCC 9595 suspension ( $10^8$  CFU/mL) was added. The cultures were incubated under shaking (120 rpm). After 48 h, the samples were serially diluted using phosphate-buffered saline (PBS) and plated on MRS agar. The plates were then incubated at 37 °C for 48 hours, and the colony-forming units (CFU) were expressed as CFU/mL<sup>9</sup>.

## **2.2. Evaluation of juice stability during storage**

### **2.2.1. Viability of *Lacticaseibacillus rhamnosus* ATCC 9595 in pasteurized Cupuaçu juice**

Samples of fermented and non-fermented juices were stored at 8 °C. After each determined period (7, 14, 21, and 28 days of refrigeration), the viability of *L. rhamnosus* ATCC 9595 was analyzed as described above.

### **2.2.2. Microbiological and physicochemical analysis**

The search for pathogenic microorganisms in the stored juices was carried out according to Normative Instruction No. 60 of 2019<sup>13</sup>. During the storage period, possible variations in color, total ash, moisture, and sugar contents were analyzed<sup>14</sup>. Five samples per batch were produced for each analysis.

## **2.3. *In vitro* simulation of the gastrointestinal tract of *Lacticaseibacillus rhamnosus* ATCC 9595 in Cupuaçu juice**

### **2.3.1. Lysozyme resistance**

Initially, the microorganisms grown in 10 mL of MRS broth at 37 °C were centrifuged, washed twice, and suspended in 2 mL with phosphate buffer (0.1 M, pH 7.0). Then, 10% of the bacterial suspension was inoculated into a sterile electrolytic solution (SEE) (0.22 g/L CaCl<sub>2</sub>, 6.2 g/L NaCl, 2.2 g/L KCl, 1.2 g/L NaHCO<sub>3</sub>) in the presence of 100 mg/L of lysozyme (Sigma-Aldrich, St. Louis, USA). For control, the bacterial suspension was also inoculated in SEE without lysozyme. The survival rate was expressed as a percentage of CFU/mL after 30 min and 120 min compared to the count determined at time zero.

**BROMATOLOGICAL ANALYSIS AND EVALUATION OF ANTIOXIDANT PROPERTIES OF  
PASTEURIZED CUPUAÇU (*Theobroma grandiflorum*) JUICE FERMENTED BY  
*Lactobacillus rhamnosus* ATCC 9595**

### **2.3.2. Bile salt resistance**

The bile salt tolerance test was performed using a previously described method, with modifications<sup>15</sup>. MRS medium and sterile non-fermented juice solutions were prepared containing bile salt (Oxgall Sigma, USA) at 0.5% or 1.0% and pH 7.3. Samples (100 µL) of each solution and 10 µL of the fermented juice were added to the wells of a 96-well plate. The plate was incubated for 3 hours at 37 °C, and the results were read at 630 nm.

### **2.4. *In vitro* antioxidant assays of cupuaçu juice**

Samples (100 mL) of fermented and non-fermented juices were subjected to liquid-liquid extraction using ethyl acetate (1:1; v/v). After separation, the solvent was removed by evaporation, and the extracts were stored at -20 °C.

#### **2.4.1. DPPH and ABTS assays**

The free radical scavenging properties of the extracts were measured using the DPPH (2,2-diphenyl-1-picrylhydrazyl; Sigma-Aldrich) radical and the ABTS (2,2'-azino-bis-(3-ethylbenzothiazoline-6-sulfonic acid) radical, according to previously described methods<sup>16</sup>.

#### **2.4.2. Determination of phenolic compounds**

The phenolic compounds were determined in the extracts using the Folin-Ciocalteu reagent. Samples (200 µL at 1000 µg/mL) were added to 1.0 mL of Folin-Ciocalteu reagent (1:1 v/v). After 3 minutes, 800 µL of sodium carbonate (20%) was added. The mixture was incubated at room temperature, protected from light, and left to stand for 2 hours. The absorbance of the mixture was measured at 765 nm. The total phenol content was expressed in µg/mL of gallic acid equivalents (GAE) using a calibration curve obtained with the standard gallic acid solution<sup>16</sup>.

### **2.5. Determination of flavonoids**

Aliquots of the samples (100 µL at different concentrations) were mixed with 100 µL of the reagent solution (2 g of aluminum chloride diluted in a 2% ethanol solution). The mixture was incubated at room temperature and protected from light, and after 60 minutes, the absorbance was measured at 420 nm. The flavonoid amount was calculated in µg/mL of

**BROMATOLOGICAL ANALYSIS AND EVALUATION OF ANTIOXIDANT PROPERTIES OF  
PASTEURIZED CUPUAÇU (*Theobroma grandiflorum*) JUICE FERMENTED BY  
*Lacticaseibacillus rhamnosus* ATCC 9595**

quercetin equivalent (QE), using a calibration curve constructed with the standard quercetin solution <sup>16</sup>.


## 2.6. Data analysis

The experiments were performed in triplicate and in three independent assays. Excel® was used for data tabulation, and GraphPad Prism version 8.1 was used for data analysis using Student's t-tests or One-way ANOVA or Two-way ANOVA, depending on the type of experiment. All results are expressed as the mean values of the groups and analyzed, considering a p-value < 0.05 as statistically significant.

## 3. RESULTS AND DISCUSSION

### 3.1. Viability of *Lacticaseibacillus rhamnosus* ATCC 9595 in pasteurized cupuaçu juice during storage

Initially, the growth of *L. rhamnosus* ATCC 9595 in pasteurized cupuaçu juice was analyzed, resulting in a population of approximately 9 Log CFU/mL, which remained stable during the 28 days of storage ( $p > 0.05$ ) (Figure 1). The juice pH after fermentation was  $4.2 \pm 0.03$ , indicating the production of organic acids. These data are consistent with those obtained in the fermentation of cupuaçu juice sterilized by autoclaving <sup>9</sup>.



**Figure 1:** Storage for 28 days of *Lacticaseibacillus rhamnosus* ATCC 9595 in pasteurized *Theobroma grandiflorum* juice.  $p > 0.05$ . Values were analyzed by One-way ANOVA test.

**BROMATOLOGICAL ANALYSIS AND EVALUATION OF ANTIOXIDANT PROPERTIES OF  
PASTEURIZED CUPUAÇU (*Theobroma grandiflorum*) JUICE FERMENTED BY  
*Lactobacillus rhamnosus* ATCC 9595**

The use of probiotics in food products in Brazil requires a prior analysis by the National Health Surveillance Agency (ANVISA), which recommends that the probiotic population should be in the range of 8 to 9 Log CFU/mL in the product ready for consumption<sup>17-19</sup>. In this context, the product formulated in this research meets the probiotic concentration required by ANVISA. In addition, it was confirmed that cupuaçu juice is a suitable matrix for the propagation of probiotic bacteria, as previously demonstrated<sup>9</sup>.

Although more commonly associated with milk, lactic fermentation can be carried out using plant matrices, such as fruit pulps and milks, as long as they contain fermentable sugars in their compositions<sup>20</sup>. The incorporation of probiotics into plant-based beverages enhances their nutritional value, intensifies or adds functional characteristics, and extends the shelf life of the beverages<sup>9,21</sup>.

Furthermore, developing fruit juices and other plant products with probiotics allows people with milk restrictions (vegans, lactose intolerant, and those with cow's milk protein allergy) to access the multiple benefits of consuming these microorganisms<sup>21,22</sup>. Recent studies demonstrate that consuming probiotics is crucial for balancing the intestinal microbiota and improving digestive and immune function<sup>23</sup>. Additionally, probiotic supplementation has beneficial effects on mental health and in treating infectious and chronic diseases<sup>24,25</sup>.

### 3.2. Microbiological analyses

Microbiological analyses revealed that the samples of pasteurized cupuaçu juice, whether fermented or not, met the standards established by Brazilian guidelines for juices<sup>13</sup>, with none of the following pathogens being detectable throughout the 28 days of storage (Table 1).

**Table 1:** Microbiological analysis of pasteurized cupuaçu juice (*Theobroma grandiflorum*).

Pathogen (/mL)	T0	T7	T14	T21	T28
	C/FC	C/FC	C/FC	C/FC	C/FC
Molds and Yeasts	0/0	0/0	0/0	0/0	0/0
<i>Enterobacteriaceae</i>	0/0	0/0	0/0	0/0	0/0
<i>Salmonella</i>	0/0	0/0	0/0	0/0	0/0
<i>Escherichia coli</i>	0/0	0/0	0/0	0/0	0/0

C = Cupuaçu juice; FC = Fermented cupuaçu juice

**BROMATOLOGICAL ANALYSIS AND EVALUATION OF ANTIOXIDANT PROPERTIES OF  
PASTEURIZED CUPUAÇU (*Theobroma grandiflorum*) JUICE FERMENTED BY  
*Lacticaseibacillus rhamnosus* ATCC 9595**

### 3.3. Physicochemical analysis

During the storage period, the possible physicochemical changes of the products were also analyzed. No significant changes were detected in the ash content, moisture, or Brix ( $p > 0.05$ ) in the analyzed periods. Regarding the pH, the non-fermented juice showed a significant increase at the end of the storage period ( $p < 0.05$ ) (Table 2).

**Table 2:** Physicochemical analysis of pasteurized cupuaçu juices (*Theobroma grandiflorum*).

	Ash (%)		Moisture (%)		Brix (B°)		pH	
	C	FC	C	FC	C	FC	C	FC
T0	0.81 ± 0.04 <sup>a,1</sup>	0.86 ± 0.06 <sup>a,1</sup>	96.92 ± 0.20 <sup>a,1</sup>	96.83 ± 0.20 <sup>a,1</sup>	3 <sup>a,1</sup>	3 <sup>a,1</sup>	5.16 ± 0.05 <sup>a,1</sup>	4.2 ± 0.03 <sup>a,2</sup>
T7	0.81 ± 0.04 <sup>a,1</sup>	0.72 ± 0.10 <sup>a,1</sup>	96.66 ± 0.03 <sup>a,1</sup>	96.61 ± 0.001 <sup>a,1</sup>	3 <sup>a,1</sup>	3 <sup>a,1</sup>	5.26 ± 0.05 <sup>b,1</sup>	4.4 ± 0.03 <sup>b,2</sup>
T14	0.78 ± 0.20 <sup>a,1</sup>	0.84 ± 0.10 <sup>a,1</sup>	96.78 ± 0.002 <sup>a,1</sup>	96.75 ± 0.03 <sup>a,1</sup>	3 <sup>a,1</sup>	3 <sup>a,1</sup>	5.12 ± 0.04 <sup>a,1</sup>	4.32 ± 0.00 <sup>c,2</sup>
T21	0.83 ± 0.20 <sup>a,1</sup>	0.83 ± 0.08 <sup>a,1</sup>	96.56 ± 0.1 <sup>a,1</sup>	96.55 ± 0.05 <sup>a,1</sup>	3 <sup>a,1</sup>	3 <sup>a,1</sup>	5.00 ± 0.01 <sup>c,1</sup>	4.27 ± 0.01 <sup>c,2</sup>
T28	0.81 ± 0.50 <sup>a,1</sup>	0.56 ± 0.20 <sup>a,1</sup>	96.71 ± 0.01 <sup>a,1</sup>	96.60 ± 0.00 <sup>a,1</sup>	3 <sup>a,1</sup>	3 <sup>a,1</sup>	5.95 ± 0.01 <sup>d,1</sup>	4.34 ± 0.02 <sup>c,2</sup>


C = Cupuaçu juice; FC = Fermented cupuaçu juice. In each column, values with statistically significant differences are indicated by the superscript letters (a, b, c) different. For each parameter analyzed at a specific time (rows), values with statistically significant differences are indicated by superscript numbers (1, 2) different. Values were analyzed by Two-way ANOVA test.

### 3.4. *In vitro* simulation of the gastrointestinal tract of *Lacticaseibacillus rhamnosus* ATCC 9595 in cupuaçu juice

Next, the survival of *L. rhamnosus* ATCC 9595 in cupuaçu juice under various conditions found in the gastrointestinal tract was evaluated (Figure 2). *L. rhamnosus* ATCC 9595 resisted the action of lysozyme (100 mg/L) when cultured in cupuaçu juice and MRS broth (Figure 2A). However, a greater resistance was observed in cupuaçu juice, at the two analyzed times ( $p < 0.05$ ). These data were similar to those reported for *L. rhamnosus* ATCC 9595 grown in bacuri juice<sup>12</sup>. Similarly, *L. rhamnosus* ATCC 9595 resisted exposure to bile salts (0.5% to 1.0%), obtaining similar variations when cultured in MRS or cupuaçu juice (Figure 2B).



**BROMATOLOGICAL ANALYSIS AND EVALUATION OF ANTIOXIDANT PROPERTIES OF  
PASTEURIZED CUPUAÇU (*Theobroma grandiflorum*) JUICE FERMENTED BY  
*Lactiseibacillus rhamnosus* ATCC 9595**




**Figure 2:** Evaluation of the tolerance to simulated gastrointestinal conditions. (A) Lysozyme tolerance test. (B) Bile salts tolerance test. \* $p < 0.05$ ; \*\*\*\* $p < 0.0001$ . Values were analyzed by Two-way ANOVA test.

### 3.5. Effect of fermentation on the antioxidant properties and levels of phenolic compounds and flavonoids in cupuaçu juice

The antioxidant capacity of cupuaçu juice, fermented or not by *L. rhamnosus* ATCC 9595, was evaluated by DPPH and ABTS assays (Figure 3). It was observed that the antioxidant properties of cupuaçu juice significantly increased due to fermentation by *L. rhamnosus* ATCC 9595. In the comparative analysis, significant differences were observed at all concentrations in the DPPH assay and at concentrations  $\leq 250 \mu\text{g/mL}$  in the ABTS assay. The fermented juice samples subjected to storage showed IC<sub>50</sub> values similar to those of day 0, while the non-fermented extract showed variation from day 21 onwards (Figure 3C).

The results obtained with the fermentation of pasteurized cupuaçu juice are consistent with several studies demonstrating the efficacy of lactic fermentation by *L. rhamnosus* strains (isolated or in co-culture) in increasing the antioxidant properties of fruit juices, such as nettle-fortified orange juice<sup>26</sup> and mixed juçara and mango juice<sup>27</sup>.


**BROMATOLOGICAL ANALYSIS AND EVALUATION OF ANTIOXIDANT PROPERTIES OF  
PASTEURIZED CUPUAÇU (*Theobroma grandiflorum*) JUICE FERMENTED BY  
*Lactiseibacillus rhamnosus* ATCC 9595**



**Figure 3:** Evaluation of the antioxidant properties of pasteurized *Theobroma grandiflorum* juices. (A) Evaluation of antioxidant activity by the DPPH method. (B) Evaluation of antioxidant activity by the ABTS method; (C) Evaluation of antioxidant activity by the DPPH method of samples obtained during storage. IC50 = concentration that inhibits 50% of the radical; \*\*\*\* p < 0.0001. Values were analyzed by Two-way ANOVA test.

Figure 4 presents the total phenolic compounds and flavonoids levels in fermented or non-fermented cupuaçu juice samples by *L. rhamnosus*. It was observed that fermentation significantly increased the levels of total phenolic compounds ( $p < 0.01$ ); however, no changes were detected in the flavonoid content.

**BROMATOLOGICAL ANALYSIS AND EVALUATION OF ANTIOXIDANT PROPERTIES OF  
PASTEURIZED CUPUAÇU (*Theobroma grandiflorum*) JUICE FERMENTED BY  
*Lactiseibacillus rhamnosus* ATCC 9595**



**Figure 4:** Measurement of total phenolic compounds (A) and flavonoids (B) in fermented and non-fermented cupuaçu juices. GAE: gallic acid equivalent; QE: quercetin equivalent. \*\*  $p < 0.01$ . Values were analyzed by the Student's t-test.

A diet rich in phenolic compounds is associated with a decreased risk of myocardial infarction, stroke, and diabetes, as it improves lipid profile, blood pressure, insulin resistance, and systemic inflammation<sup>28</sup>. Particularly, flavonoids are abundant in fruits and responsible for their color, flavor, and aroma. Due to their antioxidant and anti-inflammatory characteristics, they are important in preventing and treating cardiovascular diseases, diabetes, neurological disorders, and cancer<sup>29</sup>.

#### 4. CONCLUSION

The results obtained in this study reaffirm the potential of cupuaçu juice in the development of probiotic beverages. The juice samples (fermented or non-fermented by *L. rhamnosus* ATCC 9595) remained in accordance with Brazilian guidelines even after 28 days of refrigeration. The stored samples of both juices showed no significant changes in ash content, moisture, or Brix values during the analyzed periods; however, an increase in pH values was detected only for the non-fermented juice. Importantly, the population of *L. rhamnosus* ATCC

**BROMATOLOGICAL ANALYSIS AND EVALUATION OF ANTIOXIDANT PROPERTIES OF  
PASTEURIZED CUPUAÇU (*Theobroma grandiflorum*) JUICE FERMENTED BY  
*Lactocaseibacillus rhamnosus* ATCC 9595**

9595 remained stable during storage. *L. rhamnosus* ATCC 9595 cultured in cupuaçu juice showed resistance to the adverse conditions simulated by the gastrointestinal tract. Additionally, fermentation with *L. rhamnosus* ATCC 9595 significantly increased the antioxidant characteristics of cupuaçu juice, a finding related to the increase in total phenolic compounds.

In this context, it is suggested that the fermentation of cupuaçu juice by *L. rhamnosus* ATCC 9595 is an efficient strategy for enhancing antioxidant properties. The beverage produced is a lactose-free probiotic product, appealing to the vegan market and individuals with intolerances and allergies. It is also a functional food that can potentially alleviate pathological damage induced by free radicals.

## REFERENCES

- 1 Zaric BL, Macvanin MT, Isenovic ER. Free radicals: Relationship to Human Diseases and Potential Therapeutic applications. *Int J Biochem Cell Biol* 2023; 154. doi:10.1016/J.BIOCEL.2022.106346.
- 2 Demirci-Çekiç S, Özkan G, Avan AN, Uzunboy S, Çapanoğlu E, Apak R. Biomarkers of Oxidative Stress and Antioxidant Defense. *J Pharm Biomed Anal* 2022; 209: 114477.
- 3 Sharifi-Rad M, Anil Kumar N v., Zucca P, Varoni EM, Dini L, Panzarini E *et al.* Lifestyle, Oxidative Stress, and Antioxidants: Back and Forth in the Pathophysiology of Chronic Diseases. *Front Physiol* 2020; 11: 694.
- 4 Matzembacher dos Santos J, Gomes Heck T. Oxidative stress in type 2 diabetes and the impact of exercise: from mitochondria to glucose management in skeletal muscle. *Revista Contexto & Saúde* 2022; 22: e11425–e11425.
- 5 Forman HJ, Zhang H. Targeting oxidative stress in disease: promise and limitations of antioxidant therapy. *Nature Reviews* 2021; 20: 689.
- 6 Giampieri F, Battino M. Bioactive Phytochemicals and Functional Food Ingredients in Fruits and Vegetables. *Int J Mol Sci* 2020; 21. doi:10.3390/IJMS21093278.
- 7 Punaro GR, Lima DY, Rodrigues AM, Pugliero S, Mouro MG, Rogero MM *et al.* Cupuaçu extract reduces nitrosative stress and modulates inflammatory mediators in the kidneys of experimental diabetes. *Clinical Nutrition* 2019; 38: 364–371.
- 8 Rodrigues DBB, Punaro GR, de LIMA DY, Rodrigues AM, Pugliero S, Higa EMS. Cupuaçu extract protects the kidneys of diabetic rats by modulating Nrf2/NF-κB p65 and iNOS. *An Acad Bras Cienc* 2023; 95: e20220927.
- 9 Zagnignan A, Mendes YC, Mesquita GP, Santos GDC dos, Silva L dos S, de Souza Sales AC *et al.* Short-Term Intake of *Theobroma grandiflorum* Juice Fermented with *Lactocaseibacillus rhamnosus*

**BROMATOLOGICAL ANALYSIS AND EVALUATION OF ANTIOXIDANT PROPERTIES OF  
PASTEURIZED CUPUAÇU (*Theobroma grandiflorum*) JUICE FERMENTED BY  
*Lactocaseibacillus rhamnosus* ATCC 9595**

- ATCC 9595 Amended the Outcome of Endotoxemia Induced by Lipopolysaccharide. *Nutrients* 2023; 15. doi:10.3390/NU15041059.
- 10 Ribeiro FC, de Barros PP, Rossoni RD, Junqueira JC, Jorge AOC. Lactobacillus rhamnosus inhibits Candida albicans virulence factors in vitro and modulates immune system in Galleria mellonella. *J Appl Microbiol* 2017; 122: 201–211.
  - 11 Mendi A, Köse S, Uçkan D, Akca G, Yilmaz D, Aral L *et al.* Lactobacillus rhamnosus could inhibit Porphyromonas gingivalis derived CXCL8 attenuation. *Journal of Applied Oral Science* 2016; 24: 67–75.
  - 12 Mendes YC, Mesquita GP, Costa GDE, Barbosa da Silva AC, Gouveia E, Silva MRC *et al.* Evaluation of Growth, Viability, Lactic Acid Production and Anti-Infective Effects of Lactocaseibacillus rhamnosus ATCC 9595 in Bacuri Juice (Platonia insignis). *Foods* 2021; 10: 603.
  - 13 Brasil. Agência Nacional de Vigilância Sanitária, INSTRUÇÃO NORMATIVA N° 60, DE 23 DE DEZEMBRO DE 2019 - INSTRUÇÃO NORMATIVA N° 60, DE 23 DE DEZEMBRO DE 2019 - DOU - Imprensa Nacional. 2019<https://www.in.gov.br/web/dou/-/instrucao-normativa-n-60-de-23-de-dezembro-de-2019-235332356> (accessed 17 Aug2023).
  - 14 Zenebon O, Pascuet NS. Métodos físico-químicos para análise de alimentos. 2005; : 1018–1018.
  - 15 Silva B, Jung LRC, Sandes SHC, Alvim LB, Bomfim MRQ, Nicoli JR *et al.* In vitro assessment of functional properties of lactic acid bacteria isolated from faecal microbiota of healthy dogs for potential use as probiotics. *Benef Microbes* 2013.
  - 16 Silva TF, Cavalcanti Filho JRN, Fonsêca MMLB, Santos NM dos, Silva ACB da, Zagnignan A *et al.* Products Derived from Buchenavia tetraphylla Leaves Have In Vitro Antioxidant Activity and Protect Tenebrio molitor Larvae against Escherichia coli-Induced Injury. *Pharmaceuticals* 2020; 13: 46.
  - 17 Brasil. Agência Nacional de Vigilância Sanitária. RDC N° 241, DE 26 DE JULHO DE 2018. 2018<http://antigo.anvisa.gov.br/legislacao#/visualizar/378665> (accessed 17 Aug2023).
  - 18 Brasil. Agência Nacional de Vigilância Sanitária. Guia para instrução processual de petição de avaliação de probióticos para uso em alimentos. 2021<http://antigo.anvisa.gov.br/guias#/visualizar/448269> (accessed 17 Aug2023).
  - 19 Brasil. Agência Nacional de Vigilância Sanitária. Alimentos com alegações de propriedades funcionais e ou de saúde, novos alimentos/ingredientes, substâncias bioativas e probióticos: IX lista de alegações de propriedade funcional aprovada, atualizado em julho/2008. 2008.
  - 20 Ruiz Rodríguez LG, Zamora Gasga VM, Pescuma M, Van Nieuwenhove C, Mozzi F, Sánchez Burgos JA. Fruits and fruit by-products as sources of bioactive compounds. Benefits and trends of lactic acid fermentation in the development of novel fruit-based functional beverages. *Food Research International* 2021; 140: 109854.
  - 21 Naseem Z, Mir SA, Wani SM, Rouf MA, Bashir I, Zehra A. Probiotic-fortified fruit juices: Health benefits, challenges, and future perspective. *Nutrition* 2023; 115. doi:10.1016/J.NUT.2023.112154.

**BROMATOLOGICAL ANALYSIS AND EVALUATION OF ANTIOXIDANT PROPERTIES OF  
PASTEURIZED CUPUAÇU (*Theobroma grandiflorum*) JUICE FERMENTED BY  
*Lactobacillus rhamnosus* ATCC 9595**

- 22 Maia MS, Domingos MM, de São José JFB. Viability of Probiotic Microorganisms and the Effect of Their Addition to Fruit and Vegetable Juices. *Microorganisms* 2023; 11. doi:10.3390/MICROORGANISMS11051335.
- 23 Dahiya D, Nigam PS. Nutrition and Health through the Use of Probiotic Strains in Fermentation to Produce Non-Dairy Functional Beverage Products Supporting Gut Microbiota. *Foods* 2022; 11. doi:10.3390/FOODS11182760.
- 24 Schneider E, Doll JPK, Schweinfurth N, Kettelhack C, Schaub AC, Yamanbaeva G *et al.* Effect of short-term, high-dose probiotic supplementation on cognition, related brain functions and BDNF in patients with depression: a secondary analysis of a randomized controlled trial. *J Psychiatry Neurosci* 2023; 48: E23–E33.
- 25 Gutiérrez-Castrellón P, Gandara-Martí T, Abreu Y Abreu AT, Nieto-Rufino CD, López-Orduña E, Jiménez-Escobar I *et al.* Probiotic improves symptomatic and viral clearance in Covid19 outpatients: a randomized, quadruple-blinded, placebo-controlled trial. *Gut Microbes* 2022; 14. doi:10.1080/19490976.2021.2018899.
- 26 Sengun IY, Kirmizigul A, Atlama K, Yilmaz B. The viability of *Lactobacillus rhamnosus* in orange juice fortified with nettle (*Urtica dioica* L.) and bioactive properties of the juice during storage. *LWT* 2020; 118: 108707.
- 27 Prates FC, de Castro Leite Júnior BR, Martins EMF, Cristianini M, da Silva RR, da Rocha Campos AN *et al.* Development of a mixed jussara and mango juice with added *Lactobacillus rhamnosus* GG submitted to sub-lethal acid and baric stresses. *J Food Sci Technol* 2020; 57: 4524–4532.
- 28 Rana A, Samtiya M, Dhewa T, Mishra V, Aluko RE. Health benefits of polyphenols: A concise review. *J Food Biochem* 2022; 46. doi:10.1111/JFBC.14264.
- 29 Dias MC, Pinto DCGA, Silva AMS, Giovinazzo G, Gerardi C, Mosca L. Plant Flavonoids: Chemical Characteristics and Biological Activity. *Molecules* 2021, Vol 26, Page 5377 2021; 26: 5377.

Submitted: August 21, 2023

Accepted: March 25, 2024

Published: May 6, 2024

**Author contributions:**

Daniel dos Santos Guimarães – Methodology, Validation, Visualization.

Camila Caetano da Silva - Methodology, Validation, Visualization, Writing – original draft.

Kátia Sayuri Aragão Aguiar - Methodology, Validation.

Alexia Figueiredo Ferreira - - Methodology, Validation.

Marcos Andrade Silva - Methodology

**BROMATOLOGICAL ANALYSIS AND EVALUATION OF ANTIOXIDANT PROPERTIES OF  
PASTEURIZED CUPUAÇU (*Theobroma grandiflorum*) JUICE FERMENTED BY  
*Lactobacillus rhamnosus* ATCC 9595**

Luís Cláudio Nascimento da Silva - Conceptualization, Formal analysis, Obtaining funding, project administration, supervision and writing of the original manuscript

Adrielle Zagnignan – Conceptualization, Formal analysis, Obtaining funding, project administration, supervision and writing of the original manuscript.

Todos os autores aprovaram a versão final do texto.

Conflict of interest: There is no conflict of interest.

This work was funded by the Maranhão Scientific and Technological Research and Development Support Foundation (Process: Universal 00881/19, POS-GRAD-02460/21, INFRA-02032/21) and Conselho Nacional de Desenvolvimento Científico e Tecnológico (Process: 312349/2020-3)

**Corresponding author:**

Adrielle Zagnignan

Universidade Ceuma

Rua Anapurus, n.1, Renascença II, São Luís/MA, Brasil - CEP: 65075-120

E-mail: [adrielle004602@ceuma.com.br](mailto:adrielle004602@ceuma.com.br)

Editor: Dr, Matias Nunes Frizzo

Editor-in-chief: Dra. Adriane Cristina Bernat Kolankiewicz

This is an open access article distributed under the terms of the license Creative Commons.

